

## Datasheet for ABIN5653497

# **CYP1B1 ELISA Kit**



Go to Product page

_					
	W	0	rv	10	W

Quantity:	96 tests
Target:	CYP1B1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	1.56 ng/mL - 100 ng/mL
Minimum Detection Limit:	1.56 ng/mL
Application:	ELISA

#### **Product Details**

Sample Type:	Cell Lysate, Tissue Homogenate	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Specificity:	This assay has high sensitivity and excellent specificity for detection of Cytochrome P450 1B1 (CYP1B1). No significant cross-reactivity or interference between Cytochrome P450 1B1 (CYP1B1) and analogues was observed.	
Sensitivity:	0.54 ng/mL	

## **Target Details**

Target:	CYP1B1
Alternative Name:	Cytochrome P450 1B1 (CYP1B1 Products)

## **Target Details**

Background:	Gene Name: Cytochrome P450 1B1		
	Gene Aliases: CP1B, GLC3A, Cytochrome P450,Family 1,Subfamily B,Polypeptide 1, Glaucoma		
	3,Primary Infantile		
Gene ID:	1545		
UniProt:	Q16678		
Pathways:	Steroid Hormone Biosynthesis		
Application Details			
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than		
	5 % within the expiration date under appropriate storage condition. To minimize extra influence		
	on the performance, operation procedures and lab conditions, especially room temperature, air		
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that		
	the whole assay is performed by the same operator from the beginning to the end.		
Assay Time:	3 h		
Plate:	Pre-coated Pre-coated		
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate		
	provided in this kit has been pre-coated with an antibody specific to Cytochrome P450 1B1		
	(CYP1B1). Standards or samples are then added to the appropriate microtiter plate wells with a		
	biotin-conjugated antibody specific to Cytochrome P450 1B1 (CYP1B1). Next, Avidin		
	conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated.		
	After TMB substrate solution is added, only those wells that contain Cytochrome P450 1B1		
	(CYP1B1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in		
	color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution		
	and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm.		
	The concentration of Cytochrome P450 1B1 (CYP1B1) in the samples is then determined by		
	comparing the O.D. of the samples to the standard curve.		
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level		
	Cytochrome P450 1B1 (CYP1B1) were tested 20 times on one plate, respectively		
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level		
	Cytochrome P450 1B1 (CYP1B1) were tested on 3 different plates, 8 replicates in each plate.		
	CV(%) = SD/meanX100		
	Intra-Assay: CV<10%		

## **Application Details**

Restrictions:	For Research Use only	
Handling		
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.	
Storage:	4 °C,-20 °C	
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.	
Expiry Date:	4-8 months	