

# Datasheet for ABIN5653530

#### **CRLF1 ELISA Kit**



#### Overview

Quantity:	96 tests
Target:	CRLF1
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	15.62 pg/mL - 1000 pg/mL
Minimum Detection Limit:	15.62 pg/mL
Application:	ELISA

#### **Product Details**

Sample Type:	Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Cytokine Receptor Like Factor 1 (CRLF1). No significant cross-reactivity or interference between Cytokine Receptor Like Factor 1 (CRLF1) and analogues was observed.
Sensitivity:	5.9 pg/mL

# Target Details

Target:	CRLF1
Alternative Name:	Cytokine Receptor Like Factor 1 (CRLF1 Products)

### **Target Details**

Background:	Gene Name: Cytokine Receptor Like Factor 1
	Gene Aliases: CISS, CISS1, CLF, CLF-1, NR6, Cytokine-like factor 1, ZcytoR5
Gene ID:	12931
UniProt:	Q9JM58
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Cytokine Receptor Like
	Factor 1 (CRLF1). Standards or samples are then added to the appropriate microtiter plate
	wells with a biotin-conjugated antibody specific to Cytokine Receptor Like Factor 1 (CRLF1).
	Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and
	incubated. After TMB substrate solution is added, only those wells that contain Cytokine
	Receptor Like Factor 1 (CRLF1), biotin-conjugated antibody and enzyme-conjugated Avidin will
	exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of
	sulphuric acid solution and the color change is measured spectrophotometrically at a
	wavelength of 450nm ± 10nm. The concentration of Cytokine Receptor Like Factor 1 (CRLF1) in
	the samples is then determined by comparing the O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Cytokine Receptor Like Factor 1 (CRLF1) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Cytokine Receptor Like Factor 1 (CRLF1) were tested on 3 different plates, 8 replicates in each
	plate. CV(%) = SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only

# Handling

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months