

Datasheet for ABIN5653583 **DEFB103A ELISA Kit**



Overview

Quantity:	96 tests
Target:	DEFB103A
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	31.25 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.25 pg/mL
Application:	ELISA

Product Details

Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Specificity:	This assay has high sensitivity and excellent specificity for detection of Defensin Beta 103B (DEFB103B). No significant cross-reactivity or interference between Defensin Beta 103B (DEFB103B) and analogues was observed.	
Sensitivity:	12.7 pg/mL	
Target Details		
Target:	DEFB103A	
Alternative Name:	Defensin Beta 103B (DEFB103A Products)	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5653583 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Background: Pathways:	Gene Name: Defensin Beta 103B Gene Aliases: HBP3, HBD3, BD3, DEFB103, DEFB3, Defensin-like protein, Beta-defensin 3 Production of Molecular Mediator of Immune Response
Pathways:	
Pathways:	Production of Molecular Mediator of Immune Response
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Defensin Beta 103B
	(DEFB103B). Standards or samples are then added to the appropriate microtiter plate wells with
	a biotin-conjugated antibody specific to Defensin Beta 103B (DEFB103B). Next, Avidin
	conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated.
	After TMB substrate solution is added, only those wells that contain Defensin Beta 103B
	(DEFB103B), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in
	color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution
	and the color change is measured spectrophotometrically at a wavelength of 450nm \pm 10nm.
	The concentration of Defensin Beta 103B (DEFB103B) in the samples is then determined by
	comparing the O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Defensin Beta 103B (DEFB103B) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Defensin Beta 103B (DEFB103B) were tested on 3 different plates, 8 replicates in each plate.
	CV(%) = SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5653583 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Н	land	lina
11	ana	mig

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months