

Datasheet for ABIN5653757 **DYNC1H1 ELISA Kit**



Overview

Quantity:	96 tests
Target:	DYNC1H1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Dynein, Cytoplasmic 1, Heavy Chain 1 (DYNC1H1). No significant cross-reactivity or interference between Dynein, Cytoplasmic 1, Heavy Chain 1 (DYNC1H1) and analogues was observed.
Sensitivity:	0.113 ng/mL
Target Details	

Target:	DYNC1H1
Alternative Name:	Dynein, Cytoplasmic 1, Heavy Chain 1 (DYNC1H1 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5653757 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Gene Name: Dynein, Cytoplasmic 1, Heavy Chain 1 Gene Aliases: DHC1, DHC1a, DNCH1, DNCL, DNECL, DYHC, Dnchc1, HL-3, p22, Dynein heavy chain, cytosolic, Cytoplasmic dynein heavy chain 1, Cytoplasmic dynein 1 heavy chain 1
Gene ID:	1778
UniProt:	Q14204
Pathways:	M Phase, Ribonucleoprotein Complex Subunit Organization

Application Details

Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Dynein, Cytoplasmic 1,
	Heavy Chain 1 (DYNC1H1). Standards or samples are then added to the appropriate microtiter
	plate wells with a biotin-conjugated antibody specific to Dynein, Cytoplasmic 1, Heavy Chain 1
	(DYNC1H1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each
	microplate well and incubated. After TMB substrate solution is added, only those wells that
	contain Dynein, Cytoplasmic 1, Heavy Chain 1 (DYNC1H1), biotin-conjugated antibody and
	enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is
	terminated by the addition of sulphuric acid solution and the color change is measured
	spectrophotometrically at a wavelength of 450nm \pm 10nm. The concentration of Dynein,
	Cytoplasmic 1, Heavy Chain 1 (DYNC1H1) in the samples is then determined by comparing the
	O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Dynein, Cytoplasmic 1, Heavy Chain 1 (DYNC1H1) were tested 20 times on one plate,
	respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Dynein, Cytoplasmic 1, Heavy Chain 1 (DYNC1H1) were tested on 3 different plates, 8 replicates
	in each plate. CV(%) = SD/meanX100

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5653757 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Application Details	
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months