

Datasheet for ABIN5653866

ECE1 ELISA Kit



_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	96 tests
Target:	ECE1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	31.25 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.25 pg/mL
Application:	ELISA

Product Details

Sample Type:	Plasma, Serum	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Specificity:	This assay has high sensitivity and excellent specificity for detection of Endothelin Converting Enzyme 1 (ECE1). No significant cross-reactivity or interference between Endothelin Converting Enzyme 1 (ECE1) and analogues was observed.	
Sensitivity:	11.8 pg/mL	

Target Details

Target:	ECE1	
Alternative Name:	Endothelin Converting Enzyme 1 (ECE1 Products)	

Target Details

Background:	Gene Name: Endothelin Converting Enzyme 1
Gene ID:	1889
UniProt:	P42892
Pathways:	Regulation of Systemic Arterial Blood Pressure by Hormones, cAMP Metabolic Process, Regulation of G-Protein Coupled Receptor Protein Signaling
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Endothelin Converting Enzyme 1 (ECE1). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Endothelin Converting Enzyme 1 (ECE1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Endothelin Converting Enzyme 1 (ECE1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Endothelin Converting Enzyme 1 (ECE1) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Endothelin Converting Enzyme 1 (ECE1) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Endothelin Converting Enzyme 1 (ECE1) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Restrictions:	For Research Use only

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn | International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com | Page 2/3 | Product datasheet for ABIN5653866 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Handling

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months