

Datasheet for ABIN5654274

FGF5 ELISA Kit



Overview

Quantity:	96 tests
Target:	FGF5
Reactivity:	Human
Method Type:	Competition ELISA
Detection Range:	12.35 pg/mL - 1000 pg/mL
Minimum Detection Limit:	12.35 pg/mL
Application:	ELISA

Product Details

Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Fibroblast Growth Factor 5 (FGF5). No significant cross-reactivity or interference between Fibroblast Growth Factor 5 (FGF5) and analogues was observed.
Sensitivity:	4.99 pg/mL

Target Details

Target:	FGF5
Alternative Name:	Fibroblast Growth Factor 5 (FGF5 Products)

Target Details

Background:	Gene Name: Fibroblast Growth Factor 5
	Gene Aliases: HBGF5, Smag82, Heparin-binding growth factor 5
Gene ID:	2250
UniProt:	Q8NF90
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, ai
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	2 h
Plate:	Pre-coated
Protocol:	This assay employs the competitive inhibition enzyme immunoassay technique. A monoclona
	antibody specific to Fibroblast Growth Factor 5 (FGF5) has been pre-coated onto a microplate
	A competitive inhibition reaction is launched between biotin labeled Fibroblast Growth Factor 9
	(FGF5) and unlabeled Fibroblast Growth Factor 5 (FGF5) (Standards or samples) with the pre-
	coated antibody specific to Fibroblast Growth Factor 5 (FGF5). After incubation the unbound
	conjugate is washed off. Next, avidin conjugated to Horseradish Peroxidase (HRP) is added to
	each microplate well and incubated. The amount of bound HRP conjugate is reverse
	proportional to the concentration of Fibroblast Growth Factor 5 (FGF5) in the sample. After
	addition of the substrate solution, the intensity of color developed is reverse proportional to the
	concentration of Fibroblast Growth Factor 5 (FGF5) in the sample.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Fibroblast Growth Factor 5 (FGF5) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Fibroblast Growth Factor 5 (FGF5) were tested on 3 different plates, 8 replicates in each plate.
	CV(%) = SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only

Handling

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months