

Datasheet for ABIN5654311 FGFRL1 ELISA Kit



Overview

Quantity:	96 tests
Target:	FGFRL1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

Product Details

Alternative Name:

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Fibroblast Growth Factor Receptor Like Protein 1 (FGFRL1). No significant cross-reactivity or interference betweer Fibroblast Growth Factor Receptor Like Protein 1 (FGFRL1) and analogues was observed.
Sensitivity:	0.051 ng/mL
Target Details	
Target:	FGFRL1

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5654311 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Fibroblast Growth Factor Receptor Like Protein 1 (FGFRL1 Products)

Target Details			
Background:	Gene Name: Fibroblast Growth Factor Receptor Like Protein 1		
	Gene Aliases: FGFR5, FHFR, FGF homologous factor receptor, FGFR-like protein, Fibroblast		
	growth factor receptor 5		
Gene ID:	53834		
UniProt:	Q8N441		
Pathways:	RTK Signaling		
Application Details			
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than		
	5 % within the expiration date under appropriate storage condition. To minimize extra influence		
	on the performance, operation procedures and lab conditions, especially room temperature, air		
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that		
	the whole assay is performed by the same operator from the beginning to the end.		
Assay Time:	3 h		
Plate:	Pre-coated		
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate		
	provided in this kit has been pre-coated with an antibody specific to Fibroblast Growth Factor		
	Receptor Like Protein 1 (FGFRL1). Standards or samples are then added to the appropriate		
	microtiter plate wells with a biotin-conjugated antibody specific to Fibroblast Growth Factor		
	Receptor Like Protein 1 (FGFRL1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is		
	added to each microplate well and incubated. After TMB substrate solution is added, only those		
	wells that contain Fibroblast Growth Factor Receptor Like Protein 1 (FGFRL1), biotin-		
	conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-		
	substrate reaction is terminated by the addition of sulphuric acid solution and the color change		
	is measured spectrophotometrically at a wavelength of 450nm \pm 10nm. The concentration of		
	Fibroblast Growth Factor Receptor Like Protein 1 (FGFRL1) in the samples is then determined		
	by comparing the O.D. of the samples to the standard curve.		
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level		
	Fibroblast Growth Factor Receptor Like Protein 1 (FGFRL1) were tested 20 times on one plate,		
	respectively		
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level		
	Fibroblast Growth Factor Receptor Like Protein 1 (FGFRL1) were tested on 3 different plates, 8		

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5654311 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

replicates in each plate. CV(%) = SD/meanX100

Application Details	
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months