

Datasheet for ABIN5655067

TIM3 ELISA Kit



Overview

Quantity:	96 tests
Target:	TIM3 (TIM 3)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	78.12 pg/mL - 5000 pg/mL
Minimum Detection Limit:	78.12 pg/mL
Application:	ELISA

Product Details

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Hepatitis A Virus Cellular Receptor 2 (HAVCR2). No significant cross-reactivity or interference between Hepatitis A Virus Cellular Receptor 2 (HAVCR2) and analogues was observed.
Sensitivity:	33 pg/mL

Target Details

Target:	TIM3 (TIM 3)
Alternative Name:	Hepatitis A Virus Cellular Receptor 2 (TIM 3 Products)

Target Details

Virus
Gene Name: Hepatitis A Virus Cellular Receptor 2 Gene Aliases: KIM-3, TIM3, TIMD3, Tim-3, T-cell membrane protein 3, T-cell immunoglobulin mucin receptor 3, T-cell immunoglobulin and mucin domain-containing protein 3
84868
Q8TDQ0
Regulation of Lipid Metabolism by PPARalpha, Cancer Immune Checkpoints
The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.
3 h
Pre-coated
The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Hepatitis A Virus Cellular Receptor 2 (HAVCR2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Hepatitis A Virus Cellular Receptor 2 (HAVCR2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Hepatitis A Virus Cellular Receptor 2 (HAVCR2), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Hepatitis A Virus Cellular Receptor 2 (HAVCR2) in the samples is then determined by comparing the O.D. o the samples to the standard curve.

Application Details

	each plate. CV(%) = SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months