

## Datasheet for ABIN5655448

# **IGFBP7 ELISA Kit**



Go to Product page

_					
	W	0	rv	10	W

Quantity:	96 tests
Target:	IGFBP7
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	125.00 pg/mL - 8000 pg/mL
Minimum Detection Limit:	125.00 pg/mL
Application:	ELISA

### **Product Details**

Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Specificity:	This assay has high sensitivity and excellent specificity for detection of Insulin Like Growth Factor Binding Protein 7 (IGFBP7). No significant cross-reactivity or interference between Insulin Like Growth Factor Binding Protein 7 (IGFBP7) and analogues was observed.	
Sensitivity:	47 pg/mL	

## Target Details

Target:	IGFBP7	
Alternative Name:	ernative Name: Insulin Like Growth Factor Binding Protein 7 (IGFBP7 Products)	

## Target Details

raiget betails		
Background:	Gene Name: Insulin Like Growth Factor Binding Protein 7  Gene Aliases: FSTL2, IGFBP-7v, MAC25, PSF, TAF, IGFBP-rP1, MAC25 protein, Prostacyclinstimulating factor, PGI2-stimulating factor, Tumor-derived adhesion factor	
Gene ID:	29817	
UniProt:	Q61581	
Pathways:	Growth Factor Binding	
Application Details		
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.	
Assay Time:	3 h	
Plate:	Pre-coated	
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Insulin Like Growth Factor Binding Protein 7 (IGFBP7). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Insulin Like Growth Factor Binding Protein 7 (IGFBP7). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Insulin Like Growth Factor Binding Protein 7 (IGFBP7), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Insulin Like Growth Factor Binding Protein 7 (IGFBP7) in the samples is then determined by comparing the O.D. of the samples to the standard curve.	
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Insulin Like Growth Factor Binding Protein 7 (IGFBP7) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Insulin Like Growth Factor Binding Protein 7 (IGFBP7) were tested on 3 different plates, 8	

# **Application Details**

	Intra-Assay: CV<10%	
	Inter-Assay: CV<12%	
Restrictions:	For Research Use only	
Handling		
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and	
	specimen samples should be assayed in duplicate. Once the procedure has been started, all	
	steps should be completed without interruption.	
Storage:	4 °C,-20 °C	
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at	
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant	
	pack. Minimize freeze/thaw cycles.	
Expiry Date:	4-8 months	