

# Datasheet for ABIN5655485

### **ITGA11 ELISA Kit**



#### Overview

Quantity:	96 tests
Target:	ITGA11
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

### **Product Details**

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Integrin Alpha 11 (ITGa11). No significant cross-reactivity or interference between Integrin Alpha 11 (ITGa11) and analogues was observed.
Sensitivity:	0.124 ng/mL

### Target Details

Target:	ITGA11
Alternative Name:	Integrin Alpha 11 (ITGA11 Products)

### Target Details

Gene ID: 22801  UniProt: Q9UKX5  Pathways: Integrin Complex  Application Details  Comment: The stability of kit is determined by the loss rate of activity. The loss rate of the stability of the expiration date under appropriate storage condition. To on the performance, operation procedures and lab conditions, especial humidity, incubator temperature should be strictly controlled. It is also the whole assay is performed by the same operator from the beginning.  Assay Time: 3 h  Plate: Pre-coated  Protocol: The test principle applied in this kit is Sandwich enzyme immunoassay provided in this kit has been pre-coated with an antibody specific to Integrin Alpha 11 (ITGa11). Next, Aviding Horseradish Peroxidase (HRP) is added to each microplate well and in	
UniProt: Q9UKX5  Pathways: Integrin Complex  Application Details  Comment: The stability of kit is determined by the loss rate of activity. The loss rate 5 % within the expiration date under appropriate storage condition. To on the performance, operation procedures and lab conditions, especial humidity, incubator temperature should be strictly controlled. It is also the whole assay is performed by the same operator from the beginning.  Assay Time: 3 h  Plate: Pre-coated  Protocol: The test principle applied in this kit is Sandwich enzyme immunoassay provided in this kit has been pre-coated with an antibody specific to Int Standards or samples are then added to the appropriate microtiter plate conjugated antibody specific to Integrin Alpha 11 (ITGa11). Next, Avidit Horseradish Peroxidase (HRP) is added to each microplate well and in	
Pathways: Integrin Complex  Application Details  Comment: The stability of kit is determined by the loss rate of activity. The loss rate 5 % within the expiration date under appropriate storage condition. To on the performance, operation procedures and lab conditions, especial humidity, incubator temperature should be strictly controlled. It is also the whole assay is performed by the same operator from the beginning.  Assay Time: 3 h  Plate: Pre-coated  Protocol: The test principle applied in this kit is Sandwich enzyme immunoassay provided in this kit has been pre-coated with an antibody specific to Integrand Standards or samples are then added to the appropriate microtiter plate conjugated antibody specific to Integrin Alpha 11 (ITGa11). Next, Aviding Horseradish Peroxidase (HRP) is added to each microplate well and in	
Application Details  Comment:  The stability of kit is determined by the loss rate of activity. The loss ra 5 % within the expiration date under appropriate storage condition. To on the performance, operation procedures and lab conditions, especial humidity, incubator temperature should be strictly controlled. It is also the whole assay is performed by the same operator from the beginning.  Assay Time:  3 h  Plate:  Pre-coated  Protocol:  The test principle applied in this kit is Sandwich enzyme immunoassay provided in this kit has been pre-coated with an antibody specific to Integrin Alpha 11 (ITGa11). Next, Aviding Horseradish Peroxidase (HRP) is added to each microplate well and incomplete well and incomplete the storage of the stable of of the	
Comment:  The stability of kit is determined by the loss rate of activity. The loss rate 5 % within the expiration date under appropriate storage condition. To on the performance, operation procedures and lab conditions, especial humidity, incubator temperature should be strictly controlled. It is also the whole assay is performed by the same operator from the beginning.  Assay Time:  3 h  Plate:  Pre-coated  Protocol:  The test principle applied in this kit is Sandwich enzyme immunoassay provided in this kit has been pre-coated with an antibody specific to Integrated Standards or samples are then added to the appropriate microtiter plated conjugated antibody specific to Integrin Alpha 11 (ITGa11). Next, Aviding Horseradish Peroxidase (HRP) is added to each microplate well and incomplete well an	
5 % within the expiration date under appropriate storage condition. To on the performance, operation procedures and lab conditions, especial humidity, incubator temperature should be strictly controlled. It is also the whole assay is performed by the same operator from the beginning.  Assay Time:  3 h  Plate:  Pre-coated  Protocol:  The test principle applied in this kit is Sandwich enzyme immunoassay provided in this kit has been pre-coated with an antibody specific to Int Standards or samples are then added to the appropriate microtiter plat conjugated antibody specific to Integrin Alpha 11 (ITGa11). Next, Aviding Horseradish Peroxidase (HRP) is added to each microplate well and incomplete well and incomplete the storage of the provided to the appropriate well and incomplete the provided to the appropriate well and the provided to the appropriate t	
Protocol:  The test principle applied in this kit is Sandwich enzyme immunoassay provided in this kit has been pre-coated with an antibody specific to Int Standards or samples are then added to the appropriate microtiter plat conjugated antibody specific to Integrin Alpha 11 (ITGa11). Next, Avidir Horseradish Peroxidase (HRP) is added to each microplate well and incomplete to the samples are the sadded to each microplate well and incomplete to the samples are the sadded to each microplate well and incomplete the samples are the sadded to each microplate well and incomplete the samples are the sadded to each microplate well and incomplete the samples are the sadded to each microplate well and incomplete the samples are the sadded to each microplate well and incomplete the samples are the sadded to each microplate well and incomplete the samples are the sadded to each microplate well and incomplete the samples are the sadded to each microplate well and the samples are the sampl	minimize extra influence ly room temperature, air strongly suggested that
Protocol: The test principle applied in this kit is Sandwich enzyme immunoassay provided in this kit has been pre-coated with an antibody specific to Int Standards or samples are then added to the appropriate microtiter plat conjugated antibody specific to Integrin Alpha 11 (ITGa11). Next, Avidin Horseradish Peroxidase (HRP) is added to each microplate well and incomplete well and in	
provided in this kit has been pre-coated with an antibody specific to Int Standards or samples are then added to the appropriate microtiter plat conjugated antibody specific to Integrin Alpha 11 (ITGa11). Next, Avidir Horseradish Peroxidase (HRP) is added to each microplate well and incomplete well and incomplete well.	
substrate solution is added, only those wells that contain Integrin Alpha conjugated antibody and enzyme-conjugated Avidin will exhibit a change substrate reaction is terminated by the addition of sulphuric acid solution is measured spectrophotometrically at a wavelength of 450nm ± 10nm Integrin Alpha 11 (ITGa11) in the samples is then determined by comparables to the standard curve.	egrin Alpha 11 (ITGa11) te wells with a biotin- n conjugated to cubated. After TMB a 11 (ITGa11), biotin- ge in color. The enzyme- on and the color change n. The concentration of
Assay Precision:  Intra-assay Precision (Precision within an assay): 3 samples with low, r Integrin Alpha 11 (ITGa11) were tested 20 times on one plate, respectiv Inter-assay Precision (Precision between assays): 3 samples with low, Integrin Alpha 11 (ITGa11) were tested on 3 different plates, 8 replicate SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%	vely middle and high level
Restrictions: For Research Use only	

## Handling

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months