

# Datasheet for ABIN5655630 **IRF1 ELISA Kit**



#### Overview

Quantity:	96 tests
Target:	IRF1
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

#### **Product Details**

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Interferon Regulatory Factor 1 (IRF1). No significant cross-reactivity or interference between Interferon Regulatory Factor 1 (IRF1) and analogues was observed.
Sensitivity:	0.056 ng/mL

## **Target Details**

Target:	IRF1
Alternative Name:	Interferon Regulatory Factor 1 (IRF1 Products)

#### **Target Details**

rarget Details	
Background:	Gene Name: Interferon Regulatory Factor 1
	Gene Aliases: MAR
Gene ID:	16362
UniProt:	P15314
Pathways:	Interferon-gamma Pathway, Response to Growth Hormone Stimulus, Positive Regulation of Immune Effector Process, Hepatitis C, Autophagy
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Interferon Regulatory Factor 1 (IRF1). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Interferon Regulatory Factor 1 (IRF1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Interferon Regulatory Factor 1 (IRF1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Interferon Regulatory Factor 1 (IRF1) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Interferon Regulatory Factor 1 (IRF1) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Interferon Regulatory Factor 1 (IRF1) were tested on 3 different plates, 8 replicates in each

plate. CV(%) = SD/meanX100

Intra-Assay: CV<10% Inter-Assay: CV<12%

## **Application Details**

Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months