

Datasheet for ABIN5655856 IL-19 CLIA Kit



Overview

| Quantity: | 96 tests |
|--------------------------|--------------------------|
| Target: | IL-19 (IL19) |
| Reactivity: | Human |
| Method Type: | Sandwich ELISA |
| Detection Range: | 15.62 pg/mL - 1000 pg/mL |
| Minimum Detection Limit: | 15.62 pg/mL |
| Application: | ELISA |

Product Details

| Sample Type: | Plasma, Serum |
|--------------------|---|
| Analytical Method: | Quantitative |
| Detection Method: | Chemiluminescent |
| Specificity: | This assay has high sensitivity and excellent specificity for detection of Interleukin 19 (IL19). No significant cross-reactivity or interference between Interleukin 19 (IL19) and analogues was observed. |
| Sensitivity: | 0.63 pg/mL |
| Target Details | |
| Target: | IL-19 (IL19) |
| Alternative Name: | Interleukin 19 (IL19 Products) |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5655856 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

| Target Details | |
|---------------------|---|
| Background: | Gene Name: Interleukin 19 Gene Aliases: MDA1, ZMDA1, IL-10C, NG.1, Melanoma Differentiation Associated Protein-Like Protein |
| Gene ID: | 29949 |
| UniProt: | Q9UHD0 |
| Application Details | |
| Comment: | The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end. |
| Assay Time: | 2 - 3 h |
| Plate: | Pre-coated |
| Protocol: | The microplate provided in this kit has been pre-coated with an antibody specific to Interleukin 19 (IL19). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Interleukin 19 (IL19). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Interleukin 19 (IL19) level in the sample or standard., |
| Assay Precision: | Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Interleukin 19 (IL19) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Interleukin 19 (IL19) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12% |
| Restrictions: | For Research Use only |
| Handling | |
| Handling Advice: | Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5655856 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

| | assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption. |
|------------------|--|
| Storage: | 4 °C,-20 °C |
| Storage Comment: | -20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles. |
| Expiry Date: | 4-8 months |