

## Datasheet for ABIN5656195 KRT4 ELISA Kit



Overview

Quantity:	96 tests
Target:	KRT4
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

## Product Details

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Keratin 4 (KRT4). No significant cross-reactivity or interference between Keratin 4 (KRT4) and analogues was observed.
Sensitivity:	0.057 ng/mL
Target Details	
Target:	KDTA

Target:	KRT4
Alternative Name:	Keratin 4 (KRT4 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5656195 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Background:	Gene Name: Keratin 4
background.	Gene Aliases: CYK4, CK4, K4, Cytokeratin 4, Keratin, type II cytoskeletal 4, Type-II keratin Kb4
UniProt:	P19013
	P19015
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Keratin 4 (KRT4).
	Standards or samples are then added to the appropriate microtiter plate wells with a biotin-
	conjugated antibody specific to Keratin 4 (KRT4). Next, Avidin conjugated to Horseradish
	Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution
	is added, only those wells that contain Keratin 4 (KRT4), biotin-conjugated antibody and
	enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is
	terminated by the addition of sulphuric acid solution and the color change is measured
	spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Keratin 4
	(KRT4) in the samples is then determined by comparing the O.D. of the samples to the standard
	curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Keratin 4 (KRT4) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Keratin 4 (KRT4) were tested on 3 different plates, 8 replicates in each plate. CV(%) =
	SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Postrictions:	
Restrictions:	For Research Use only

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5656195 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Н	land	lina
11	ana	mig

Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months