

Datasheet for ABIN5656322

PSMB9 ELISA Kit



Go to Product page

()	11/	\sim	r\	ń	0	۱۸/	,
\cup	V	C	rv	ı	ヒ	٧V	

Quantity:	96 tests
Target:	PSMB9
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

Product Details

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Large Multifunctional Peptidase 2 (LMP2). No significant cross-reactivity or interference between Large Multifunctional Peptidase 2 (LMP2) and analogues was observed.
Sensitivity:	0.051 ng/mL

Target Details

Target:	PSMB9
Alternative Name:	Large Multifunctional Peptidase 2 (PSMB9 Products)

Target Details

Background:	Gene Name: Large Multifunctional Peptidase 2		
	Gene Aliases: PSMB9, Proteasome(Prosome,Macropain)Subunit Beta Type 9, Macropain chain		
	7, Multicatalytic endopeptidase complex chain 7, Really interesting new gene 12 protein		
Gene ID:	5698		
UniProt:	P28065		
Pathways:	Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA		
Application Details			
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than		
	5 % within the expiration date under appropriate storage condition. To minimize extra influence		
	on the performance, operation procedures and lab conditions, especially room temperature, air		
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that		
	the whole assay is performed by the same operator from the beginning to the end.		
Assay Time:	3 h		
Plate:	Pre-coated		
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate		
	provided in this kit has been pre-coated with an antibody specific to Large Multifunctional		
	Peptidase 2 (LMP2). Standards or samples are then added to the appropriate microtiter plate		
	wells with a biotin-conjugated antibody specific to Large Multifunctional Peptidase 2 (LMP2).		
	Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and		
	incubated. After TMB substrate solution is added, only those wells that contain Large		
	Multifunctional Peptidase 2 (LMP2), biotin-conjugated antibody and enzyme-conjugated Avidin		
	will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of		
	sulphuric acid solution and the color change is measured spectrophotometrically at a		
	wavelength of 450nm ± 10nm. The concentration of Large Multifunctional Peptidase 2 (LMP2)		
	in the samples is then determined by comparing the O.D. of the samples to the standard curve		
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level		
	Large Multifunctional Peptidase 2 (LMP2) were tested 20 times on one plate, respectively		
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level		
	Large Multifunctional Peptidase 2 (LMP2) were tested on 3 different plates, 8 replicates in each		
	plate. CV(%) = SD/meanX100		
	Intra-Assay: CV<10%		
	Inter-Assay: CV<12%		

Application Details

Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months