

# Datasheet for ABIN5656410

# **LILRA3 ELISA Kit**



#### Overview

Quantity:	96 tests
Target:	LILRA3
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

### **Product Details**

Sample Type:	Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 (LILRA3). No significant cross-reactivity or interference between Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 (LILRA3) and analogues was observed.
Sensitivity:	0.124 ng/mL

# Target Details

Target:	LILRA3
Alternative Name:	Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 (LILRA3 Products)

# **Target Details** Gene Name: Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 Background: Gene Aliases: CD85e, HM31, HM43, ILT6, LIR4, E3, CD85 antigen-like family member E, Immunoglobulin-like transcript 6, Leukocyte immunoglobulin-like receptor 4 Gene ID: 11026 UniProt: **Q8N6C8 Application Details** Comment: The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end. Assay Time: 3 h Pre-coated Plate: Protocol:

The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 (LILRA3). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 (LILRA3). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 (LILRA3), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm  $\pm$  10nm. The concentration of Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 (LILRA3) in the samples is then determined by comparing the 0.D. of the samples to the standard curve.

Assay Precision: Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 (LILRA3) were tested 20
times on one plate, respectively
Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
Leukocyte Immunoglobulin Like Receptor Subfamily A, Member 3 (LILRA3) were tested on 3
different plates, 8 replicates in each plate. CV(%) = SD/meanX100

# **Application Details**

	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months