

# Datasheet for ABIN5656494

#### **LRP2 ELISA Kit**



_			
( )	V/C	rv	٨/

Quantity:	96 tests
Target:	LRP2
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

#### **Product Details**

Sample Type:	Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Low Density Lipoprotein Receptor Related Protein 2 (LRP2). No significant cross-reactivity or interference between Low Density Lipoprotein Receptor Related Protein 2 (LRP2) and analogues was observed.
Sensitivity:	0.067 ng/mL

# Target Details

Target:	LRP2
Alternative Name: Low Density Lipoprotein Receptor Related Protein 2 (LRP2 Products)	

#### **Target Details**

Target Details		
Background:	Gene Name: Low Density Lipoprotein Receptor Related Protein 2  Gene Aliases: DBS, gp330, Glycoprotein 330, Megalin, Low Density Lipoprotein-Related Protein	
Gene ID:	4036	
UniProt:	P98164	
Pathways:	Metabolism of Steroid Hormones and Vitamin D, Thyroid Hormone Synthesis, Hormone Transport	
Application Details		
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.	
Assay Time:	3 h	
Plate:	Pre-coated	
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Low Density Lipoprotein Receptor Related Protein 2 (LRP2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Low Density Lipoprotein Receptor Related Protein 2 (LRP2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Low Density Lipoprotein Receptor Related Protein 2 (LRP2), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Low Density Lipoprotein Receptor Related Protein 2 (LRP2) in the samples is then determined by comparing the O.D. of the samples to the standard curve.	
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Lov Density Lipoprotein Receptor Related Protein 2 (LRP2) were tested 20 times on one plate,	

replicates in each plate. CV(%) = SD/meanX100

Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level

Low Density Lipoprotein Receptor Related Protein 2 (LRP2) were tested on 3 different plates, 8

respectively

# **Application Details**

	Intra-Assay: CV<10% Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months