

Datasheet for ABIN5656887

MAP1LC3A ELISA Kit



Go	+~	$\bigcap \nu$	\sim	110+	no	00
(¬()	1()	м	()(1	11 1 () 1	1)2	$(1 \leftarrow$

_				
	۱۱ / ۱	rv		۱۸/
	' V '	 ı v	Ι.	v v

Quantity:	96 tests
Target:	MAP1LC3A
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	1.56 ng/mL - 100 ng/mL
Minimum Detection Limit:	1.56 ng/mL
Application:	ELISA

Product Details

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Microtubule Associated Protein 1 Light Chain 3 Alpha (MAP1LC3a). No significant cross-reactivity or interference between Microtubule Associated Protein 1 Light Chain 3 Alpha (MAP1LC3a) and analogues was observed.
Sensitivity:	0.59 ng/mL

Target Details

Target:	MAP1LC3A
Alternative Name:	Microtubule Associated Protein 1 Light Chain 3 Alpha (MAP1LC3A Products)

Target Details

rarget betails		
Background:	Gene Name: Microtubule Associated Protein 1 Light Chain 3 Alpha	
	Gene Aliases: MAP1ALC3, MAP1BLC3, LC3, LC3A, ATG8E, Autophagy-related ubiquitin-like	
	modifier LC3 A, MAP1 light chain 3-like protein 1	
Gene ID:	84557	
UniProt:	Q9H492	
Pathways:	Autophagy	
Application Details		
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than	
	5% within the expiration date under appropriate storage condition. To minimize extra influence	
	on the performance, operation procedures and lab conditions, especially room temperature, air	
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that	
	the whole assay is performed by the same operator from the beginning to the end.	
Assay Time:	3 h	
Plate:	Pre-coated	
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate	
	provided in this kit has been pre-coated with an antibody specific to Microtubule Associated	
	Protein 1 Light Chain 3 Alpha (MAP1LC3a). Standards or samples are then added to the	
	appropriate microtiter plate wells with a biotin-conjugated antibody specific to Microtubule	
	Associated Protein 1 Light Chain 3 Alpha (MAP1LC3a). Next, Avidin conjugated to Horseradish	
	Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution	
	is added, only those wells that contain Microtubule Associated Protein 1 Light Chain 3 Alpha	
	(MAP1LC3a), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in	
	color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution	
	and the color change is measured spectrophotometrically at a wavelength of 450nm \pm 10nm.	
	The concentration of Microtubule Associated Protein 1 Light Chain 3 Alpha (MAP1LC3a) in the	
	samples is then determined by comparing the O.D. of the samples to the standard curve.	
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level	
	Microtubule Associated Protein 1 Light Chain 3 Alpha (MAP1LC3a) were tested 20 times on one	
	plate, respectively	
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level	
	Microtubule Associated Protein 1 Light Chain 3 Alpha (MAP1LC3a) were tested on 3 different	
	plates, 8 replicates in each plate. CV(%) = SD/meanX100	

Application Details

	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months