

Datasheet for ABIN5657085

MYH1 ELISA Kit



_					
	W	0	rv	10	W

Quantity:	96 tests	
Target:	MYH1	
Reactivity:	Human	
Method Type:	Sandwich ELISA	
Detection Range:	0.156 ng/mL - 10 ng/mL	
Minimum Detection Limit:	0.156 ng/mL	
Application:	ELISA	

Product Details

Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Specificity:	This assay has high sensitivity and excellent specificity for detection of Myosin Heavy Chain 1, Skeletal Muscle, Adult (MYH1). No significant cross-reactivity or interference between Myosin Heavy Chain 1, Skeletal Muscle, Adult (MYH1) and analogues was observed.	
Sensitivity:	0.059 ng/mL	

Target Details

Target:	MYH1
Alternative Name:	Myosin Heavy Chain 1, Skeletal Muscle, Adult (MYH1 Products)

Target Details

larget Details	
Background:	Gene Name: Myosin Heavy Chain 1, Skeletal Muscle, Adult
	Gene Aliases: MYHSA1, MYHa, MyHC-2X/D, Myosin-1, Myosin Heavy Chain IIx/d
Gene ID:	4619
UniProt:	P12882
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Myosin Heavy Chain 1,
	Skeletal Muscle, Adult (MYH1). Standards or samples are then added to the appropriate
	microtiter plate wells with a biotin-conjugated antibody specific to Myosin Heavy Chain 1,
	Skeletal Muscle, Adult (MYH1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is
	added to each microplate well and incubated. After TMB substrate solution is added, only thos
	wells that contain Myosin Heavy Chain 1, Skeletal Muscle, Adult (MYH1), biotin-conjugated
	antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate
	reaction is terminated by the addition of sulphuric acid solution and the color change is
	measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of
	Myosin Heavy Chain 1, Skeletal Muscle, Adult (MYH1) in the samples is then determined by
	comparing the O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Myosin Heavy Chain 1, Skeletal Muscle, Adult (MYH1) were tested 20 times on one plate,
	respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Myosin Heavy Chain 1, Skeletal Muscle, Adult (MYH1) were tested on 3 different plates, 8
	replicates in each plate. CV(%) = SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%

Application Details

Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months