

Datasheet for ABIN5657200 **NEIL1 ELISA Kit**



Overview

Quantity:	96 tests
Target:	NEIL1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

Product Details

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Nei Endonuclease VIII Like Protein 1 (NEIL1). No significant cross-reactivity or interference between Nei Endonuclease VIII Like Protein 1 (NEIL1) and analogues was observed.
Sensitivity:	0.062 ng/mL
Target Details	
Target:	NEIL1
Alternative Name:	Nei Endonuclease VIII Like Protein 1 (NEIL1 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5657200 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Gene Name: Nei Endonuclease VIII Like Protein 1
	Gene Aliases: NEI1, hFPG1, FPG1, DNA-(apurinic or apyrimidinic site) lyase Neil1, Nei-like
	protein 1, DNA glycosylase/AP lyase Neil1
Gene ID:	79661
UniProt:	Q96FI4
Pathways:	DNA Damage Repair
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Nei Endonuclease VIII Like
	Protein 1 (NEIL1). Standards or samples are then added to the appropriate microtiter plate
	wells with a biotin-conjugated antibody specific to Nei Endonuclease VIII Like Protein 1 (NEIL1)
	Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and
	incubated. After TMB substrate solution is added, only those wells that contain Nei
	Endonuclease VIII Like Protein 1 (NEIL1), biotin-conjugated antibody and enzyme-conjugated
	Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the
	addition of sulphuric acid solution and the color change is measured spectrophotometrically at
	a wavelength of 450nm \pm 10nm. The concentration of Nei Endonuclease VIII Like Protein 1
	(NEIL1) in the samples is then determined by comparing the O.D. of the samples to the
	standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Ne
	Endonuclease VIII Like Protein 1 (NEIL1) were tested 20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Ne
	Endonuclease VIII Like Protein 1 (NEIL1) were tested on 3 different plates, 8 replicates in each
	plate. CV(%) = SD/meanX100

Intra-Assay: CV<10%

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5657200 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Application Details	
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months