

Datasheet for ABIN5657332

Neuroserpin ELISA Kit



Overview

Quantity:	96 tests
Target:	Neuroserpin (SERPINI1)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.781 ng/mL - 50 ng/mL
Minimum Detection Limit:	0.781 ng/mL
Application:	ELISA

Product Details

Sample Type:	Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Neuroserpin (NSP). No significant cross-reactivity or interference between Neuroserpin (NSP) and analogues was observed.
Sensitivity:	0.34 ng/mL

Target Details

Target:	Neuroserpin (SERPINI1)
Alternative Name:	Neuroserpin (SERPINI1 Products)

Target Details

Background:	Gene Name: Neuroserpin	
	Gene Aliases: SERPINI1, PI12, Serpin Peptidase Inhibitor Clade I Member 1, Peptidase inhibitor	
	12	
Gene ID:	5274	
UniProt:	Q99574	
Pathways:	Regulation of Hormone Metabolic Process	
Application Details		
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than	
	5% within the expiration date under appropriate storage condition. To minimize extra influence	
	on the performance, operation procedures and lab conditions, especially room temperature, air	
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that	
	the whole assay is performed by the same operator from the beginning to the end.	
Assay Time:	3 h	
Plate:	Pre-coated	
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate	
	provided in this kit has been pre-coated with an antibody specific to Neuroserpin (NSP).	
	Standards or samples are then added to the appropriate microtiter plate wells with a biotin-	
	conjugated antibody specific to Neuroserpin (NSP). Next, Avidin conjugated to Horseradish	
	Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution	
	is added, only those wells that contain Neuroserpin (NSP), biotin-conjugated antibody and	
	enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is	
	terminated by the addition of sulphuric acid solution and the color change is measured	
	spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Neuroserpin	
	(NSP) in the samples is then determined by comparing the O.D. of the samples to the standard	
	curve.	
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level	
	Neuroserpin (NSP) were tested 20 times on one plate, respectively	
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level	
	Neuroserpin (NSP) were tested on 3 different plates, 8 replicates in each plate. CV(%) =	
	SD/meanX100	
	Intra-Assay: CV<10%	
	Inter-Assay: CV<12%	

Application Details

Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months