

Datasheet for ABIN5657458 NUMA1 ELISA Kit



Overview

Quantity:	96 tests
Target:	NUMA1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

Product Details

Sample Type:	Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Nuclear Mitotic Apparatus Protein 1 (NUMA1). No significant cross-reactivity or interference between Nuclear Mitotic Apparatus Protein 1 (NUMA1) and analogues was observed.
Sensitivity:	0.057 ng/mL
Target Details	
Taraati	

Target:	NUMA1
Alternative Name:	Nuclear Mitotic Apparatus Protein 1 (NUMA1 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5657458 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Background:	Gene Name: Nuclear Mitotic Apparatus Protein 1
	Gene Aliases: NMP22, NUMA, Nuclear matrix protein-22, SP-H antigen
Gene ID:	4926
UniProt:	Q14980
Pathways:	Caspase Cascade in Apoptosis, Regulation of Actin Filament Polymerization, M Phase
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Nuclear Mitotic Apparatus
	Protein 1 (NUMA1). Standards or samples are then added to the appropriate microtiter plate
	wells with a biotin-conjugated antibody specific to Nuclear Mitotic Apparatus Protein 1
	(NUMA1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each
	microplate well and incubated. After TMB substrate solution is added, only those wells that
	contain Nuclear Mitotic Apparatus Protein 1 (NUMA1), biotin-conjugated antibody and enzyme-
	conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by
	the addition of sulphuric acid solution and the color change is measured
	spectrophotometrically at a wavelength of 450nm \pm 10nm. The concentration of Nuclear
	Mitotic Apparatus Protein 1 (NUMA1) in the samples is then determined by comparing the O.D.
	of the samples to the standard curve.
Assay Precision:	
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Nuclear Mitotic Apparatus Protein 1 (NUMA1) were tested 20 times on one plate, respectively

Nuclear Mitotic Apparatus Protein 1 (NUMA1) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100

Intra-Assay: CV<10%

Inter-Assay: CV<12%

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5657458 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Application Details	
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months