

Datasheet for ABIN5657971

PABPC1L ELISA Kit



Overview

Quantity:	96 tests
Target:	PABPC1L
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Poly A Binding Protein Cytoplasmic 1 Like Protein (PABPC1L). No significant cross-reactivity or interference between Poly A Binding Protein Cytoplasmic 1 Like Protein (PABPC1L) and analogues was observed.
Sensitivity:	0.115 ng/mL

Target Details

Target:	PABPC1L
Alternative Name:	Poly A Binding Protein Cytoplasmic 1 Like Protein (PABPC1L Products)

Target Details Gene Name: Poly A Binding Protein Cytoplasmic 1 Like Protein Background: Gene Aliases: C20orf119, PABPC1L1, ePAB, Polyadenylate-Binding Protein 1-Like Gene ID: 80336 UniProt: Q4VXU2 **Application Details** The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than Comment: 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end. Assay Time: 3 h Plate: Pre-coated Protocol: The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Poly A Binding Protein Cytoplasmic 1 Like Protein (PABPC1L). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Poly A Binding Protein Cytoplasmic 1 Like Protein (PABPC1L). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Poly A Binding Protein Cytoplasmic 1 Like Protein (PABPC1L), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Poly A Binding Protein Cytoplasmic 1 Like Protein (PABPC1L) in the samples is then determined by comparing the O.D. of the samples to the standard curve. Assay Precision:

Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Poly A Binding Protein Cytoplasmic 1 Like Protein (PABPC1L) were tested 20 times on one plate, respectively

Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Poly A Binding Protein Cytoplasmic 1 Like Protein (PABPC1L) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100

Intra-Assay: CV<10% Inter-Assay: CV<12%

Application Details

Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months