

Datasheet for ABIN5658723

SPINK1 ELISA Kit



Overview

Quantity:	96 tests
Target:	SPINK1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	1.56 ng/mL - 100 ng/mL
Minimum Detection Limit:	1.56 ng/mL
Application:	ELISA

Product Details

Sample Type:	Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Serine Peptidase Inhibitor Kazal Type 1 (SPINK1). No significant cross-reactivity or interference between Serine Peptidase Inhibitor Kazal Type 1 (SPINK1) and analogues was observed.
Sensitivity:	0.63 ng/mL

Target Details

Target:	SPINK1
Alternative Name:	Serine Peptidase Inhibitor Kazal Type 1 (SPINK1 Products)

Target Details Gene Name: Serine Peptidase Inhibitor Kazal Type 1 Background: Gene Aliases: PCTT, PSTI, TATI, Tumor-Associated Trypsin Inhibitor, Pancreatic secretory trypsin inhibitor, Tumor-associated trypsin inhibitor Gene ID: 6690 UniProt: P00995 **Application Details** Comment: The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end. Assay Time: 3 h Pre-coated Plate: Protocol: The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Serine Peptidase Inhibitor Kazal Type 1 (SPINK1). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Serine Peptidase Inhibitor Kazal Type 1 (SPINK1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Serine Peptidase Inhibitor Kazal Type 1 (SPINK1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Serine Peptidase Inhibitor Kazal Type 1 (SPINK1) in the samples is then determined by comparing the

Assay Precision: Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Serine Peptidase Inhibitor Kazal Type 1 (SPINK1) were tested 20 times on one plate, respectively

Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Serine Peptidase Inhibitor Kazal Type 1 (SPINK1) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100

O.D. of the samples to the standard curve.

Intra-Assay: CV<10%

Application Details

	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months