

# Datasheet for ABIN5658855

## **SLC1A5 ELISA Kit**



#### Overview

Quantity:	96 tests
Target:	SLC1A5
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

### **Product Details**

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Solute Carrier Family 1, Member 5 (SLC1A5). No significant cross-reactivity or interference between Solute Carrier Family 1, Member 5 (SLC1A5) and analogues was observed.
Sensitivity:	0.124 ng/mL

### **Target Details**

Target:	SLC1A5
Alternative Name:	Solute Carrier Family 1, Member 5 (SLC1A5 Products)

### **Target Details**

rai get betaile		
Background:	Gene Name: Solute Carrier Family 1, Member 5  Gene Aliases: SLC1A5, AAAT, ASCT2, ATBO, M7V1, M7VS1, R16, RDRC, Neutral Amino Acid  Transporter B(0), Baboon M7 virus receptor, Sodium-dependent neutral amino acid transporter type 2	
Gene ID:	6510	
UniProt:	Q15758	
Pathways:	Dicarboxylic Acid Transport, Warburg Effect	
Application Details		
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.	
Assay Time:	3 h	
Plate:	Pre-coated	
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Solute Carrier Family 1, Member 5 (SLC1A5). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Solute Carrier Family 1, Member 5 (SLC1A5). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Solute Carrier Family 1, Member 5 (SLC1A5), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Solute Carrier Family 1, Member 5 (SLC1A5) in the samples is then determined by comparing the O.D. of the samples to the standard curve.	
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Solute Carrier Family 1, Member 5 (SLC1A5) were tested 20 times on one plate, respectively Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Solute Carrier Family 1, Member 5 (SLC1A5) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100	

### **Application Details**

	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months