

Datasheet for ABIN5659093

CD81 ELISA Kit



Overview

Quantity:	96 tests
Target:	CD81
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Target Of The Antiproliferative Antibody 1 (TAPA1). No significant cross-reactivity or interference between Target Of The Antiproliferative Antibody 1 (TAPA1) and analogues was observed.
Sensitivity:	0.115 ng/mL

Target Details

Target:	CD81
Alternative Name:	Target Of The Antiproliferative Antibody 1 (CD81 Products)

Target Details

rarget betails	
Background:	Gene Name: Target Of The Antiproliferative Antibody 1
	Gene Aliases: CD81, S5.7, TAPA1, TSPAN28, Tetraspanin 28, Target Of Antiproliferative
	Antibody 1, 26 kDa cell surface protein TAPA-1
Gene ID:	975
UniProt:	P60033
Pathways:	Inositol Metabolic Process, Hepatitis C
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Target Of The
	Antiproliferative Antibody 1 (TAPA1). Standards or samples are then added to the appropriate
	microtiter plate wells with a biotin-conjugated antibody specific to Target Of The
	Antiproliferative Antibody 1 (TAPA1). Next, Avidin conjugated to Horseradish Peroxidase (HRP)
	is added to each microplate well and incubated. After TMB substrate solution is added, only
	those wells that contain Target Of The Antiproliferative Antibody 1 (TAPA1), biotin-conjugated
	antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate
	reaction is terminated by the addition of sulphuric acid solution and the color change is
	measured spectrophotometrically at a wavelength of 450nm \pm 10nm. The concentration of
	Target Of The Antiproliferative Antibody 1 (TAPA1) in the samples is then determined by
	comparing the O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Target Of The Antiproliferative Antibody 1 (TAPA1) were tested 20 times on one plate,
	respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Target Of The Antiproliferative Antibody 1 (TAPA1) were tested on 3 different plates, 8
	replicates in each plate. CV(%) = SD/meanX100

Application Details

	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months