

Datasheet for ABIN5659693

TP53BP1 ELISA Kit



Overview

Quantity:	96 tests
Target:	TP53BP1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA

Product Details

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Tumor Protein p53 Binding Protein 1 (TP53BP1). No significant cross-reactivity or interference between Tumor Protein p53 Binding Protein 1 (TP53BP1) and analogues was observed.
Sensitivity:	0.138 ng/mL

Target Details

Target:	TP53BP1
Alternative Name:	Tumor Protein p53 Binding Protein 1 (TP53BP1 Products)

Target Details

Background:	Gene Name: Tumor Protein p53 Binding Protein 1 Gene Aliases: 53BP1, p53BP1, p202
Gene ID:	7158
UniProt:	Q12888
Pathways:	DNA Damage Repair
Application Details	

_			
(,)	mr	$n \cap r$	١+٠
Co	1111	HEL	IL.

The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5 % within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

Assay Time:

3 h

Plate:

Pre-coated

Protocol:

The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Tumor Protein p53 Binding Protein 1 (TP53BP1). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Tumor Protein p53 Binding Protein 1 (TP53BP1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Tumor Protein p53 Binding Protein 1 (TP53BP1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Tumor Protein p53 Binding Protein 1 (TP53BP1) in the samples is then determined by comparing the 0.D. of the samples to the standard curve.

Assay Precision:

Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
Tumor Protein p53 Binding Protein 1 (TP53BP1) were tested 20 times on one plate, respectively
Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
Tumor Protein p53 Binding Protein 1 (TP53BP1) were tested on 3 different plates, 8 replicates
in each plate. CV(%) = SD/meanX100
Intra-Assay: CV<10%
Inter-Assay: CV<12%

Application Details

Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and
	specimen samples should be assayed in duplicate. Once the procedure has been started, all
	steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at
	4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant
	pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months