

Datasheet for ABIN5659717 TIE1 ELISA Kit



Overview

Quantity:	96 tests
Target:	TIE1
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	0.625 ng/mL - 40 ng/mL
Minimum Detection Limit:	0.625 ng/mL
Application:	ELISA

Product Details

Sample Type:	Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1). No significant cross-reactivity or interference between Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1) and analogues was observed.
Sensitivity:	0.255 ng/mL

Target Details

Target:	TIE1
Alternative Name:	Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (TIE1 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5659717 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Gene Name: Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 Gene Aliases: JTK14, Tyrosine Kinase With Immunoglobulin And Epidermal Growth Factor Homology Domains 1, Tyrosine-protein kinase receptor Tie-1
Gene ID:	21846
UniProt:	Q06806
Pathways:	RTK Signaling
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 % within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, air
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.

Assay Time:	3 h
Plate:	Pre-coated
Protocol:	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate
	provided in this kit has been pre-coated with an antibody specific to Tyrosine Kinase With
	Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1). Standards or samples are then
	added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to
	Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1). Next, Avidin
	conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated.
	After TMB substrate solution is added, only those wells that contain Tyrosine Kinase With
	Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1), biotin-conjugated antibody and
	enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is
	terminated by the addition of sulphuric acid solution and the color change is measured
	spectrophotometrically at a wavelength of 450nm \pm 10nm. The concentration of Tyrosine
	Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1) in the samples is then
	determined by comparing the O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1) were tested
	20 times on one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1) were tested

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5659717 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Application Details	
	on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	The Stop Solution is acidic. Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months