

Datasheet for ABIN5660019 WNT5A CLIA Kit



Overview

Quantity:	96 tests
Target:	WNT5A
Reactivity:	Human, Mouse, Rat
Method Type:	Sandwich ELISA
Detection Range:	156.25 pg/mL - 10000 pg/mL
Minimum Detection Limit:	156.25 pg/mL
Application:	ELISA

Product Details

Sample Type:	Cell Culture Supernatant, Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Chemiluminescent
Specificity:	This assay has high sensitivity and excellent specificity for detection of Wingless Type MMTV Integration Site Family, Member 5A (WNT5A). No significant cross-reactivity or interference between Wingless Type MMTV Integration Site Family, Member 5A (WNT5A) and analogues was observed.
Sensitivity:	5.1 pg/mL
Target Details	

Target:	WNT5A
Alternative Name:	Wingless Type MMTV Integration Site Family, Member 5A (WNT5A Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5660019 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Gene Name: Wingless Type MMTV Integration Site Family, Member 5A
Pathways:	WNT Signaling, Cellular Response to Molecule of Bacterial Origin, Positive Regulation of
	Immune Effector Process, Production of Molecular Mediator of Immune Response, Regulation
	of Cell Size, Tube Formation
Application Details	
Comment:	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than
	5 $\%$ within the expiration date under appropriate storage condition. To minimize extra influence
	on the performance, operation procedures and lab conditions, especially room temperature, ai
	humidity, incubator temperature should be strictly controlled. It is also strongly suggested that
	the whole assay is performed by the same operator from the beginning to the end.
Assay Time:	2 - 3 h
Plate:	Pre-coated
Protocol:	The microplate provided in this kit has been pre-coated with an antibody specific to Wingless
	Type MMTV Integration Site Family, Member 5A (WNT5A). Standards or samples are then
	added to the appropriate microplate wells with a biotin-conjugated antibody specific to
	Wingless Type MMTV Integration Site Family, Member 5A (WNT5A). Next, Avidin conjugated to
	Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the
	mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate
	development, the intensity of the emitted light is proportional to the Wingless Type MMTV
	Integration Site Family, Member 5A (WNT5A) level in the sample or standard.,
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	Wingless Type MMTV Integration Site Family, Member 5A (WNT5A) were tested 20 times on
	one plate, respectively
	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level
	Wingless Type MMTV Integration Site Family, Member 5A (WNT5A) were tested on 3 different
	plates, 8 replicates in each plate. CV(%) = SD/meanX100
	Intra-Assay: CV<10%
	Inter-Assay: CV<12%
Restrictions:	For Research Use only
Handling	
Handling Advice:	Do not allow to contact skin or eyes. Calibrators, controls and specimen samples should be

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5660019 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

	assayed in duplicate. Once the procedure has been started, all steps should be completed without interruption.
Storage:	4 °C,-20 °C
Storage Comment:	-20°C. Bring all reagents to room temperature before beginning test. The kit may be stored at 4°C for immediate use within two days upon arrival. Reseal any unused strips with desiccant pack. Minimize freeze/thaw cycles.
Expiry Date:	4-8 months