# antibodies -online.com





# anti-CD57 antibody (PE)





#### Overview

Quantity:	100 tests
Target:	CD57 (B3GAT1)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD57 antibody is conjugated to PE
Application:	Flow Cytometry (FACS)

## **Product Details**


Immunogen:	A pool of neuroblastoma cell lines
Clone:	TB01
Isotype:	IgM
Specificity:	The mouse monoclonal antibody TB01 recognizes CD57, a carbohydrate extracellular antigen present mainly on NK cells, NK T cells, and in neural tissue.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions.  Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

## **Target Details**

Target:	CD57 (B3GAT1)


# **Target Details**

rarget Details	
Alternative Name:	CD57 (B3GAT1 Products)
Background:	Beta-1,3-glucuronyltransferase 1,CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycolipids and chondroitin sulphate proteoglycans in the nervous system. It serves as a NK cell marker and it is expressed on well differentiated prostate cancers and uveal and cutaneous melanoma. CD57+ T cells are implicated as suppressors of T-cell responses.,HNK1, LEU7, 3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc
Gene ID:	27087
UniProt:	Q9P2W7
Pathways:	Glycosaminoglycan Metabolic Process
Application Details	
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 $\mu$ L reagent / 100 $\mu$ L of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
Comment:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Restrictions:	For Research Use only
Handling	
Buffer:	Stabilizing Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.




## Flow Cytometry

**Image 1.** Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD57 (TB01) PE antibody (10  $\mu$ L reagent / 100  $\mu$ L of peripheral whole blood).



#### **Flow Cytometry**

**Image 2.** Surface staining of human peripheral blood with anti-human CD57 (TB01) PE.



#### **Flow Cytometry**

**Image 3.** Flow cytometry multicolor surface staining of human lymphocytes stained using anti-human CD57 (TB01) PE antibody (10  $\mu$ L reagent / 100  $\mu$ L of peripheral whole blood) and anti-human CD8 (MEM-31) Pacific Blue antibody (4  $\mu$ L reagent / 100  $\mu$ L of peripheral whole blood).

Please check the product details page for more images. Overall 5 images are available for ABIN5665755.