

Datasheet for ABIN5668086

Recombinant anti-p53 antibody (AA 371-380)



[Go to Product page](#)

2 Images

Overview

Quantity:	200 µg
Target:	p53 (TP53)
Binding Specificity:	AA 371-380
Reactivity:	Human, Mouse, Rat, Monkey, Rabbit
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Chimeric
Conjugate:	This p53 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunoprecipitation (IP), Immunohistochemistry (Frozen Sections) (IHC (fro)), In vivo Studies (in vivo)

Product Details

Purpose:	Anti-p53 [PAb421], Rabbit IgG, kappa
Immunogen:	Synthetic peptide corresponding to aa 371-380 of human p53.
Clone:	PAb421
Isotype:	IgG kappa
Specificity:	Recognises the mammalian mutant and wild-type p53 protein and reacts with an epitope between amino acid residues 370 and 378 (near the C-terminus of the protein), corresponding to the sequence "KKGQSTRHK".

Product Details

Cross-Reactivity: Monkey, Mouse, Rabbit, Rat

Characteristics: Original Species of Ab: Mouse
Original Format of Ab: IgG2a

Purification: Protein A affinity purified

Target Details

Target: p53 (TP53)

Alternative Name: p53 (TP53 Products)

Background: Tumor suppressor p53, Cellular tumor antigen p53

UniProt: [P04637](#)

Pathways: [p53 Signaling](#), [MAPK Signaling](#), [PI3K-Akt Signaling](#), [Apoptosis](#), [AMPK Signaling](#), [Chromatin Binding](#), [ER-Nucleus Signaling](#), [Positive Regulation of Endopeptidase Activity](#), [Hepatitis C](#), [Protein targeting to Nucleus](#), [Autophagy](#), [Warburg Effect](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Comment: This chimeric rabbit antibody was made using the variable domain sequences of the original murine IgG2a format, for improved compatibility with existing reagents, assays and techniques.

Restrictions: For Research Use only

Handling

Concentration: 1 mg/mL


Buffer: PBS with 0.02 % Proclin 300.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.



Western Blotting

Image 1. Western Blot using anti-p53 antibody PAb42. A431 cell nuclear (A) and cytoplasmic (B) extract (35 µg protein in RIPA buffer) was resolved on a 10 % SDS PAGE gel and blots probed with the chimeric rabbit version of PAb421 (ABIN5668086) at 0.1 µg/mL before detection by an anti-rabbit secondary antibody. A primary incubation of 1h was used and protein was detected by chemiluminescence. The expected band size for p53 is 43.7 kDa, though due to the high number of proline residues in this protein runs at a size of ~53 kDa (c.f. Ziemer et al., PMID: 7107651). ABIN5668086 successfully detected both human nuclear and cytoplasmic p53.



Immunofluorescence

Image 2. Immunofluorescence staining of fixed A431 with anti-p53 antibody PAb421. Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15 % Triton and stained with the chimeric rabbit IgG version of PAb421 (ABIN5668086) at 10 µg/mL for 1h followed by Alexa Fluor®488 secondary antibody (1 µg/mL), showing nuclear staining. The nuclear stain is DAPI (blue). Panels show from left-right, top-bottom ABIN5668086, DAPI, merged channels and a negative control. The negative control was stained with unimmunized rabbit IgG followed by Alexa Fluor®488 secondary antibody.