

Datasheet for ABIN5674603

CSF1R Protein (AA 20-512) (His tag,AVI tag,Biotin)



[Go to Product page](#)

2 Images

Overview

Quantity:	200 µg
Target:	CSF1R
Protein Characteristics:	AA 20-512
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CSF1R protein is labelled with His tag,AVI tag,Biotin.

Product Details

Brand:	PrecisionAvi
Sequence:	AA 20-512
Specificity:	Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.
Characteristics:	This protein carries an Avi tag (Avitag™) at the C-terminus, followed by a polyhistidine tag. The protein has a calculated MW of 57.1 kDa. The protein migrates as 65-92 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

Target Details

Target:	CSF1R
Alternative Name:	M-CSF R (CSF1R Products)
Background:	Colony stimulating factor 1 receptor (CSF1R) is also known as macrophage colony-stimulating factor receptor (M-CSFR), CD115 Cluster of Differentiation 115 (CD115), C-FMS, CSFR, FIM2, FMS, and is a member of the typeIII subfamily of receptor tyrosine kinases (RTKs). CSF1R is a receptor for a cytokine called colony stimulating factor 1, The protein encoded by the CSFR1 gene is the receptor for colony stimulating factor 1, a cytokine which controls the production, differentiation, and function of macrophages. This receptor mediates most, if not all, of the biological effects of this cytokine. Ligand binding activates CSFR1 through a process of oligomerization and transphosphorylation . Mutations in CSF1R are associated with chronic myelomonocytic leukemia and type M4 acute myeloblastic leukemia. Increased levels of CSF1R1 are found in microglia in Alzheimer's disease and after brain injuries. The increased receptor expression causes microglia to become more active. Both CSF1R, and its ligand colony stimulating factor 1 play an important role in the development of the mammary gland and may be involved in the process of mammary gland carcinogenesis.
Molecular Weight:	57.1 kDa
NCBI Accession:	NP_005202
Pathways:	RTK Signaling , Inositol Metabolic Process , Cell-Cell Junction Organization


Application Details

Comment:	Ready-to-use AvitagTM biotinylated protein: The product is exclusively produced using the AvitagTM technology. Briefly, a unique 15 amino acid peptide, the Avi tag, is introduced into the recombinant protein during expression vector construction. The single lysine residue in the Avi tag is enzymatically biotinylated by the E. Coli biotin ligase BirA. This single-point enzymatic labeling technique brings many advantages for commonly used binding assays. The biotinylation happens on the lysine residue of Avi tag, and therefore does NOT interfere with the target protein's natural binding activities. In addition, when immobilized on an avidin-coated surface, the protein orientation is uniform because the position of the Avi tag in the protein is precisely controlled.
Restrictions:	For Research Use only

Handling


Format:	Lyophilized
Buffer:	PBS, pH 7.4
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C

Images



SDS-PAGE

Image 1. Biotinylated Human M-CSF R, Avitag,His Tag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95 % .



ELISA

Image 2. Immobilized Human M-CSF, His Tag (ABIN5674639,ABIN6253718) at 5 µg/mL (100 µL/well) can bind Biotinylated Human M-CSF R, Avitag,His Tag (ABIN5674603,ABIN6253703) with a linear range of 16-500 ng/mL (QC tested).