

Datasheet for ABIN5674720  
**Podocan ELISA Kit**



[Go to Product page](#)

## Overview

Quantity:	96 tests
Target:	Podocan (PODN)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	78.125 pg/mL - 5000 pg/mL
Minimum Detection Limit:	78.125 pg/mL
Application:	ELISA

## Product Details

Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Sensitivity:	26.5 pg/mL
Components:	Pre-coated, ready to use 96-well strip plate Standard (freeze dried) Standard Diluent Detection Reagent A Detection Reagent B Assay Diluent A Assay Diluent B TMB Substhumane

## Product Details

---

Stop Solution  
Wash Buffer(30 x concenthumane)  
Plate sealer for 96 wells  
Instruction manual

---

Material not included:

1. Microplate reader with  $450 \pm 10\text{nm}$  filter.
2. Precision single or multi-channel pipettes and disposable tips.
3. Eppendorf Tubes for diluting samples.
4. Deionized or distilled water.
5. Absorbent paper for blotting the microtiter plate.
6. Container for Wash Solution.

## Target Details

---

Target: Podocan (PODN)

---

Abstract: [PODN Products](#)

---

Background: Alternative name: PCAN, SLRR5A

---

Gene ID: 127435

---

UniProt: [Q7Z5L7](#)

## Application Details

---

Application Notes: Optimal working dilution should be determined by the investigator.

---

Assay Time: 1 - 4.5 h

---

Plate: Pre-coated

---

Protocol:

1. Prepare all reagents, samples and standards
2. Add 100 $\mu\text{L}$  standard or sample to each well. Incubate 2 hours at 37°C
3. Aspirate and add 100 $\mu\text{L}$  prepared Detection Reagent A. Incubate 1 hour at 37°C
4. Aspirate and wash 3 times
5. Add 100 $\mu\text{L}$  prepared Detection Reagent B. Incubate 1 hour at 37°C
6. Aspirate and wash 5 times
7. Add 90 $\mu\text{L}$  Substrate Solution. Incubate 15-25 minutes at 37°C
8. Add 50 $\mu\text{L}$  Stop Solution. Read at 450nm immediately.

---

Restrictions: For Research Use only

## Handling

---

Storage: 4 °C

---

Expiry Date: 12 months