

Datasheet for ABIN5674867
PYCARD ELISA Kit



[Go to Product page](#)

Overview

Quantity:	96 tests
Target:	PYCARD
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA

Product Details

Sample Type:	Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Sensitivity:	0065 ng/mL
Components:	Pre-coated, ready to use 96-well strip plate Standard (freeze dried) Standard Diluent Detection Reagent A Detection Reagent B Assay Diluent A Assay Diluent B TMB Substhumane

Product Details

Stop Solution
Wash Buffer(30 x concentrated)
Plate sealer for 96 wells
Instruction manual

Material not included:

1. Microplate reader with 450 ± 10 nm filter.
2. Precision single or multi-channel pipettes and disposable tips.
3. Eppendorf Tubes for diluting samples.
4. Deionized or distilled water.
5. Absorbent paper for blotting the microtiter plate.
6. Container for Wash Solution.

Target Details

Target: PYCARD

Alternative Name: PYD And CARD Domain Containing Protein (PYCARD) ([PYCARD Products](#))

Background: Alternative name: ASC, CARD5, TMS1, Apoptosis-associated speck-like protein containing a CARD, Caspase recruitment domain-containing protein 5, Target of methylation-induced silencing 1

Gene ID: 29108

UniProt: [Q9ULZ3](#)

Pathways: [Activation of Innate immune Response](#), [Cellular Response to Molecule of Bacterial Origin](#), [Regulation of Actin Filament Polymerization](#), [Positive Regulation of Endopeptidase Activity](#), [Activated T Cell Proliferation](#), [Inflammasome](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Assay Time: 1 - 4.5 h

Plate: Pre-coated

Protocol:

1. Prepare all reagents, samples and standards
2. Add 100 μ L standard or sample to each well. Incubate 2 hours at 37°C
3. Aspirate and add 100 μ L prepared Detection Reagent A. Incubate 1 hour at 37°C
4. Aspirate and wash 3 times
5. Add 100 μ L prepared Detection Reagent B. Incubate 1 hour at 37°C

Application Details

6. Aspirate and wash 5 times
 7. Add 90 μ L Substrate Solution. Incubate 15-25 minutes at 37°C
 8. Add 50 μ L Stop Solution. Read at 450nm immediately.
-

Restrictions: For Research Use only

Handling

Storage: 4 °C

Expiry Date: 12 months
