Datasheet for ABIN5674920

PDE3B ELISA Kit

Go to Product page

Overview

Quantity:	96 tests
Target:	PDE3B
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	78.125 pg/mL - 5000 pg/mL
Minimum Detection Limit:	78.125 pg/mL
Application:	ELISA

Product Details

Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Sensitivity:	26.35 pg/mL
Components:	Pre-coated, ready to use 96-well strip plate Standard (freeze dried) Standard Diluent Detection Reagent A Detection Reagent B Assay Diluent A Assay Diluent B TMB Substhumane

Copyright antibodies-online. All rights reserved.

Product Details

	Stop Solution Wash Buffer(30 x concenthumane) Plate sealer for 96 wells Instruction manual
Material not included:	1. Microplate reader with 450 ± 10nm filter. 2. Precision single or multi-channel pipettes and disposable tips. 3. Eppendorf Tubes for diluting samples. 4. Deionized or distilled water. 5. Absorbent paper for blotting the microtiter plate. 6. Container for Wash Solution.

Target Details

Target:	PDE3B
Alternative Name:	Phosphodiesterase 3B, cGMP Inhibited (PDE3B) (PDE3B Products)
Background:	Alternative name: HcGIP1, cGIPDE1, CGIPDE1, CGIP1, CGI-PDE B, Cyclic GMP-inhibited phosphodiesterase B
Gene ID:	5140
UniProt:	Q13370
Pathways:	Negative Regulation of Hormone Secretion, Carbohydrate Homeostasis, cAMP Metabolic Process

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Assay Time:	1 - 4.5 h
Plate:	Pre-coated
Protocol:	1. Prepare all reagents, samples and standards 2. Add 100µL standard or sample to each well. Incubate 2 hours at 37℃ 3. Aspirate and add 100µL prepared Detection Reagent A. Incubate 1 hour at 37℃ 4. Aspirate and wash 3 times 5. Add 100µL prepared Detection Reagent B. Incubate 1 hour at 37℃ 6. Aspirate and wash 5 times 7. Add 90µL Substrate Solution. Incubate 15-25 minutes at 37℃

Copyright antibodies-online. All rights reserved.

Application Details

	8. Add 50µL Stop Solution. Read at 450nm immediately.
Restrictions:	For Research Use only

Handling

Storage:	4 °C
Expiry Date:	12 months

 Copyright antibodies-online. All rights reserved.