antibodies -online.com





anti-UMPS antibody (AA 351-450) (Biotin)



()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	100 μL	
Target:	UMPS	
Binding Specificity:	AA 351-450	
Reactivity:	Human, Cynomolgus, Rhesus Monkey	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This UMPS antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human UMPS
Isotype:	IgG
Cross-Reactivity:	Cynomolgus, Human, Rhesus Monkey
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	UMPS
Alternative Name:	UMPS (UMPS Products)

Target Details

Background:	Synonyms: Uridine 5'-monophosphate synthase, UMPS, UMP synthase, Orotate	
	phosphoribosyltransferase, OPRT, OPRTase, ODC, OMPdecase, OK/SW-cl.21	
	Background: This gene encodes a uridine 5'-monophosphate synthase. The encoded protein is	
	a bifunctional enzyme that catalyzes the final two steps of the de novo pyrimidine biosynthetic	
	pathway. The first reaction is carried out by the N-terminal enzyme orotate	
	phosphoribosyltransferase which converts orotic acid to orotidine-5'-monophosphate. The	
	terminal reaction is carried out by the C-terminal enzyme OMP decarboxylase which converts	
	orotidine-5'-monophosphate to uridine monophosphate. Defects in this gene are the cause of	
	hereditary orotic aciduria. Alternate splicing results in multiple transcript variants.	
Gene ID:	7372	
UniProt:	P11172	
Pathways:	Ribonucleoside Biosynthetic Process	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	