antibodies -online.com





anti-ERO1LB antibody (AA 1-100) (Biotin)



Go to Product page

()	11/0	K\ /	iew
	\cup	'I V/I	$I \cap VV$

Quantity:	100 μL	
Target:	ER01LB	
Binding Specificity:	AA 1-100	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ERO1LB antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ERO1LB
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Pig,Rabbit,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	ER01LB
Alternative Name:	ER01LB (ER01LB Products)

Target Details

Background:	Synonyms: Endoplasmic oxidoreductin-1-like protein B, ERO1-L-beta, ERO1B, ERO1-like protein	
	beta, Oxidoreductin-1-L-beta	
	Background: Oxidoreductase involved in disulfide bond formation in the endoplasmic reticulum.	
	Efficiently reoxidizes P4HB/PDI, the enzyme catalyzing protein disulfide formation, in order to	
	allow P4HB to sustain additional rounds of disulfide formation. Other protein disulfide	
	isomerase family members can also be reoxidized, but at lower rates compared to P4HB,	
	including PDIA2 (50 % of P4HB reoxidation rate), as well as PDIA3, PDIA4, PDIA6 and NXNDC12	
	(<10 %). Following P4HB reoxidation, passes its electrons to molecular oxygen via FAD, leading	
	to the production of reactive oxygen species (ROS) in the cell. May be involved in oxidative	
	proinsulin folding in pancreatic cells, hence may play a role in glucose homeostasis.	
Gene ID:	56605	
UniProt:	Q86YB8	
Pathways:	SARS-CoV-2 Protein Interactome	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	