# antibodies -online.com





# anti-CPY antibody



( )	11/0	K\ /	iew
	$\cup$	'I V/I	$I \cap VV$

Overview		
Quantity:	10 mg	
Target:	CPY	
Reactivity:	Saccharomyces cerevisiae	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CPY antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Enzyme Immunoassay (EIA), Dot Blot (DB), Immunodiffusion (ID), Radioimmunoassay (RIA)	
Product Details		
Immunogen:	Carboxypeptidase Y isolated and purified from Baker's yeast. Freund's complete adjuvant is used in the first step of the immunization procedure.	
Isotype:	IgG	
Specificity:	The antibody recognizes Carboxypeptidase Y from Baker's Yeast. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques:  Immunoelectrophoresis, Cross-Immunoelectrophoresis, single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and Enzyme Inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined.	
Purification:	Ammonium Sulphate Precipitation and Ion Exchange Chromatography	

#### **Target Details**

Target:	CPY	
Alternative Name:	Carboxypeptidase Y (CPY Products)	
Background:	Carboxypeptidase Y is involved in degradation of small peptides. Digests preferentially peptides containing an aliphatic or hydrophobic residue in P1' position, as well as methionine, leucine or phenylalanine in P1 position of ester substrate. Synonyms: Carboxypeptidase YSCY, PRC1	
Gene ID:	855343	
NCBI Accession:	NP_014026	
UniProt:	P00729	

## **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	

## Handling

Reconstitution:	Restore by adding 1.0 mL of sterile distilled water	
Concentration:	10.0 mg/mL	
Buffer:	PBS, pH 7.2 without preservatives and foreign proteins	
Preservative:	Without preservative	
Storage:	4 °C/-20 °C	
Storage Comment:	Store the antibody lyophilized at 2-8 °C and reconstituted at 2-8 °C for one week or (in aliquots) at -20 °C for longer. If a slight precipitation occurs upon storage, this should be removed by centrifugation.	