

[Go to Product page](#)

Datasheet for ABIN568217 **anti-AGLU antibody (Biotin)**

Overview

| | |
|--------------|---|
| Quantity: | 1 mL |
| Target: | AGLU |
| Reactivity: | Saccharomyces cerevisiae |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This AGLU antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), Immunoprecipitation (IP), Immunofluorescence (IF), Immunodiffusion (ID), Enzyme Immunoassay (EIA), Dot Blot (DB), Radioimmunoassay (RIA) |

Product Details

| | |
|------------------|--|
| Immunogen: | Alpha-Glucosidase isolated and purified from Baker's Yeast. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Isotype: | IgG |
| Specificity: | This antibody recognizes Alpha-Glucosidase from Baker's Yeast. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined. |
| Characteristics: | Molar Ratio: Biotin/ IgG ~6.2 |
| Purification: | Ammonium Sulphate Precipitation and Ion Exchange Chromatography |

Target Details

| | |
|-------------------|---|
| Target: | AGLU |
| Alternative Name: | alpha-Glucosidase (AGLU Products) |
| Background: | Synonyms: YIL172C |
| Gene ID: | 853235 |
| NCBI Accession: | NP_012096 |
| UniProt: | P40439 |
| Pathways: | Cellular Glucan Metabolic Process |

Application Details

| | |
|--------------------|--|
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|---|
| Format: | Liquid |
| Reconstitution: | Restore by adding 1.0 mL of sterile distilled water |
| Concentration: | 10 mg/mL |
| Buffer: | PBS, pH 7.2 without preservatives and foreign proteins |
| Preservative: | Without preservative |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store the antibody lyophilized at 2-8 °C and reconstituted at 2-8 °C for one week or (in aliquots) at -20 °C for longer. If a slight precipitation occurs upon storage, this should be removed by centrifugation. |