

#### Datasheet for ABIN568237

# anti-Glycerol Kinase antibody



_					
	W	0	rv	10	W

Overview		
Quantity:	10 mg	
Target:	Glycerol Kinase (GK)	
Reactivity:	Bacillus	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Glycerol Kinase antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Enzyme Immunoassay (EIA), Immunodiffusion (ID), Radioimmunoassay (RIA), Dot Blot (DB)	
Product Details		
Immunogen:	Glycerokinase isolated and purified from Bacillus stearothermophilus. Freund's complete	
	adjuvant is used in the first step of the immunization procedure.	
Isotype:	IgG	
Purification:	Ammonium Sulphate Precipitation and Ion Exchange Chromatography	
Target Details		
Target:	Glycerol Kinase (GK)	
Alternative Name:	Glycerol Kinase (GK Products)	
Background:	Glycerol kinase catalyzes the formation of glycerol 3 phosphate from ATP and glycerol.  Dihydroxyacetone and L glyceraldehyde can also act as acceptors, UTP and, in the case of the yeast enzyme, ITP and GTP can act as donors. It provides a way for glycerol derived from fats	

#### **Target Details**

	or glycerides to enter the glycolytic pathway. Synonyms: ATP: glycerol 3-phosphotransferase, GK, GKD, Glycerokinase
Gene ID:	939741
NCBI Accession:	NP_388810
UniProt:	P18157

## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

### Handling

Reconstitution:	Restore by adding 1.0 mL of sterile distilled water
Concentration:	10.0 mg/mL
Buffer:	PBS, pH 7.2 without preservatives and foreign proteins
Preservative:	Without preservative
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody lyophilized at 2-8 °C and reconstituted at 2-8 °C for one week or (in aliquots) at -20 °C for longer. If a slight precipitation occurs upon storage, this should be removed by centrifugation.