## antibodies -online.com





## anti-Synaptophysin 2 antibody (AA 257-261)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Overview		
Quantity:	0.1 mg	
Target:	Synaptophysin 2	
Binding Specificity:	AA 257-261	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Synaptophysin 2 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	synaptophysin 2 antibody was raised against a peptide sequence around aa.257~261 (G-P-T-S-	
	F) derived from Rat synaptophysin 2.	
Specificity:	This antibody detects endogenous levels of total synaptophysin 2 protein.	
Purification:	Antibodies were purified by affinity-chromatography using epitope-specific peptide.	
Target Details		
Target:	Synaptophysin 2	
Abstract:	Synaptophysin 2 Products	
Background:	Synaptophysin (SYP) is a neuronal synaptic vesicle glycoprotein that is expressed in	
	neuroendocrine cells and neoplasms, and thus can be used as a marker in tumor diagnosis (1).	
	Synaptophysin contains four transmembrane domains that form a hexameric channel or gap	

## Target Details

junction-like pore (2). Synaptophysin binds to the SNARE protein synaptobrevin/VAMP, which
prevents the inclusion of synaptobrevin in the synaptic vesicle fusion complex and creates a
pool of synaptobrevin for exocytosis when synapse activity increases (3). Synaptophysin is also
responsible for targeting synaptobrevin 2/VAMP2 to synaptic vesicles, a critical component of
the fusion complex (4).
37 kDa

Molecular Weight:

Gene ID: 66030

NCBI Accession: NP\_076464

UniProt: P22831

## **Application Details**

Application Notes:	Western Blot: 1:500~1:1000
Restrictions:	For Research Use only

Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Antibody supplied in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store antibody at -20°C for up to one year.