

Datasheet for ABIN5690018

anti-IQSEC1 antibody (AA 60-110)



Go to Product page

| Overview | |
|----------------------|--|
| Quantity: | 0.1 mg |
| Target: | IQSEC1 |
| Binding Specificity: | AA 60-110 |
| Reactivity: | Mouse, Human, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This IQSEC1 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF) |
| Product Details | |
| Immunogen: | IQSEC1 antibody was raised against a 19 amino acid peptide near the amino terminus of |
| | human IQSEC1. |
| | The immunogen is located within amino acids 60 - 110 of IQSEC1. |
| Isotype: | IgG |
| Specificity: | IQSEC1 antibody is human, mouse and rat reactive. At least four isoforms of IQSEC1 are known |
| | to exist. IQSEC1 antibody is predicted to not cross-react with IQSEC2. |
| Purification: | IQSEC1 antibody is affinity chromatography purified via peptide column. |
| Target Details | |
| Target: | IQSEC1 |

Target Details

| Alternative Name: | IQSEC1 (IQSEC1 Products) |
|---------------------|--|
| Background: | ADP ribosylation factors (Arfs) are small GTP-binding proteins known for their role in vesicular |
| | transport, where they nucleate the assembly of coat protein complexes at sites of carrier |
| | vesicle formation. IQSEC1 (GEP100/BRAG2), a guanine nucleotide exchange protein for ARF6, |
| | belongs to the BRAG family and contains one IQ domain, one PH domain and one SEC7 domain |
| | (1). As a known activator of ARF6, IQSEC1 regulates cell surface levels of Beta1 integrin and its |
| | knockdown blocked tumour metastasis (2,3). IQSEC1 links EGFR signaling to ARF6 activation |
| | and may contribute to the cancer metastasis and malignancy (4). |
| Molecular Weight: | Predicted: 106 kDa |
| | Observed: 103 kDa |
| Gene ID: | 9922 |
| NCBI Accession: | NP_055684 |
| UniProt: | Q6DN90 |
| Application Details | |
| Application Notes: | IQSEC1 antibody can be used for detection of IQSEC1 by Western blot at 1 - 2 μ,g/mL. Antibody |
| | can also be used for immunocytochemistry at 2 $\mu\text{,g/mL}.$ For immunofluorescence start at 2 $\mu\text{,}$ |
| | g/mL. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 mg/mL |
| Buffer: | IQSEC1 antibody is supplied in PBS containing 0.02 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |
| | should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | IQSEC1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. |