

Datasheet for ABIN5690064 anti-MYH8 antibody (AA 60-110)



o to Froduct page

\sim			
()\	/er	1/16	1///
\cup	101	VIC	, V V

Quantity:	0.1 mg	
Target:	MYH8	
Binding Specificity:	AA 60-110	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MYH8 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF)	
Product Details		
Immunogen:	MYH8 antibody was raised against a 19 amino acid peptide near the amino terminus of human MYH8.	
	The immunogen is located within amino acids 60 - 110 of MYH8.	
lsotype:	lgG	
Specificity:	MYH8 antibody is human, mouse and rat reactive. MYH8 antibody is predicted to not cross- react with other members of the myosin heavy chain family.	
Purification:	MYH8 antibody is affinity chromatography purified via peptide column.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN5690064 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Target:	MYH8	
Alternative Name:	MYH8 (MYH8 Products)	
Background:	Myosins are actin-based motor proteins that function in the generation of mechanical force in eukaryotic cells (1). MYH8 (myosin, heavy chain 8, skeletal muscle, perinatal) is a member of the class II or conventional myosin heavy chains, and functions in skeletal muscle contraction (2,3). This gene is predominantly expressed in fetal skeletal muscle. MYH8 is regulated by phosphorylation via myosin light chain kinase (MLCK) and by intracellular Ca2+ concentrations (3). A mutation in this gene results in trismus-pseudocamptodactyly syndrome (4).	
Molecular Weight:	Predicted: 213 kDa Observed: 220 kDa	
Gene ID:	4626	
NCBI Accession:	NP_002463	
UniProt:	P13535	
Application Details		
Application Notes:	MYH8 antibody can be used for detection of MYH8 by Western blot at 1 - 2 μ,g/mL. Antibody can also be used for immunohistochemistry starting at 5 μ,g/mL. For immunofluorescence start at 20 μ,g/mL.	
Restrictions: Handling	For Research Use only	
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	MYH8 antibody is supplied in PBS containing 0.02 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	MYH8 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN5690064 | 07/25/2024 | Copyright antibodies-online. All rights reserved.