

Datasheet for ABIN5690749

## C-JUN ELISA Kit

### 2 Images



[Go to Product page](#)

### Overview

Quantity:	96 tests
Target:	C-JUN (JUN)
Reactivity:	Human
Method Type:	DNA-Binding ELISA
Application:	ELISA

### Product Details

Purpose:	Human c-Jun Transcription Factor Activity Assay. This assay uses a dsDNA coated plate with canonical c-Jun binding sequences to semi-quantitatively detect active c-Jun in lysates or nuclear extracts.
Sample Type:	Cell Lysate, Nuclear Extract
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The oligonucleotide/antibody pair provided in this kit recognizes mouse c-Jun in whole lysates and nuclear extracts.
Characteristics:	<ul style="list-style-type: none"><li>• Specific transcription factor-DNA binding assay</li><li>• Perfect alternative to EMSA</li><li>• Easy to perform in an ELISA format</li><li>• Non-radioactive assay</li><li>• High throughput (96 well plate format)</li><li>• Assay can be completed within 5 hours</li></ul>

## Product Details

---

Components:	<ul style="list-style-type: none"><li>• 96-well Strip Microplate pre-coated with DNA probes</li><li>• DNA Binding Buffer</li><li>• Positive Control Sample</li><li>• Specific Competitor DNA probe</li><li>• Non-specific Competitor DNA probe</li><li>• Assay Reagent</li><li>• DTT</li><li>• Wash Buffer</li><li>• Primary Antibody</li><li>• HRP-conjugated Secondary Antibody</li><li>• TMB One-Step Substrate Reagent</li><li>• Stop Solution</li></ul>
-------------	--

Material not included:	<ul style="list-style-type: none"><li>• Distilled or deionized water</li><li>• 100 mL and 1 liter graduated cylinders</li><li>• Tubes to prepare sample dilutions Absorbent paper</li><li>• Precision pipettes to deliver 2 µL to 1 mL volumes</li><li>• Adjustable 1-25 mL pipettes for reagent preparation</li><li>• Benchtop rocker or shaker</li><li>• Microplate reader capable of measuring absorbance at 450 nm</li></ul>
------------------------	--

## Target Details

---

Target:	C-JUN (JUN)
Alternative Name:	c-Jun ( <a href="#">JUN Products</a> )
Gene ID:	3725
UniProt:	<a href="#">P05412</a>
Pathways:	<a href="#">MAPK Signaling</a> , <a href="#">RTK Signaling</a> , <a href="#">WNT Signaling</a> , <a href="#">Fc-epsilon Receptor Signaling Pathway</a> , <a href="#">Activation of Innate immune Response</a> , <a href="#">Myometrial Relaxation and Contraction</a> , <a href="#">Skeletal Muscle Fiber Development</a> , <a href="#">Protein targeting to Nucleus</a> , <a href="#">Toll-Like Receptors Cascades</a> , <a href="#">Autophagy</a> , <a href="#">Signaling of Hepatocyte Growth Factor Receptor</a> , <a href="#">BCR Signaling</a> , <a href="#">S100 Proteins</a>

## Application Details

---

Plate:	Pre-coated
Protocol:	<ol style="list-style-type: none"><li>1. Prepare all reagents and samples as instructed in the manual.</li><li>2. Add 100 µL of sample or positive control to each well.</li><li>3. Incubate 2 h at RT or O/N at 4 °C.</li><li>4. Add 100 µL of prepared primary antibody to each well.</li></ol>

## Application Details

5. Incubate 1 h at RT.
6. Add 100  $\mu$ L of prepared HRP-secondary antibody to each well.
7. Incubate 1 h at RT.
8. Add 100  $\mu$ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50  $\mu$ L of Stop Solution to each well.
11. Read at 450 nm immediately.

Restrictions: For Research Use only


## Handling

Storage: 4 °C

Storage Comment: Upon receipt, the positive control should be removed and stored at -20° or -80°C. The remainder of the kit can be stored for up to 6 months at 2-8°C from the date of shipment. Opened Microplate Wells or reagents may be stored for up to 1 month at 2° to 8°C. Return unused wells to the pouch containing desiccant pack, reseal along entire edge. Note: The kit can be used within one year if the whole kit is stored at -20°C upon receipt. Avoid repeated freeze-thaw cycles.


Expiry Date: 6 months

## Images



### Activity Assay

**Image 1.** Transcription factor assay of c-Jun from nuclear extracts of K562 cells or K562 cells treated with PMA (50 ng/ml) for 3 hr. A. Western-blot result of c-Jun from cytoplasmic and nuclear fractions. B. Transcription factor assay of c-Jun from nuclear fractions with the TF Activity Assay Kit.



### Activity Assay

**Image 2.** Transcription factor assay of c-Jun from nuclear extracts of K562 cells or K562 cells treated with PMA (50 ng/ml) for 3 hr with the specific competitor or non-specific competitor. The result shows specific binding of c-Jun to the conserved binding site detected by using the c-Jun TF Activity Assay Kit.