

Datasheet for ABIN5690749

C-JUN ELISA Kit**2** Images[Go to Product page](#)

Overview

Quantity:	96 tests
Target:	C-JUN (JUN)
Reactivity:	Human
Method Type:	DNA-Binding ELISA
Application:	ELISA

Product Details

Purpose:	Human c-Jun Transcription Factor Activity Assay. This assay uses a dsDNA coated plate with canonical c-Jun binding sequences to semi-quantitatively detect active c-Jun in lysates or nuclear extracts.
Sample Type:	Cell Lysate, Nuclear Extract
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The oligonucleotide/antibody pair provided in this kit recognizes mouse c-Jun in whole lysates and nuclear extracts.
Characteristics:	<ul style="list-style-type: none">• Specific transcription factor-DNA binding assay• Perfect alternative to EMSA• Easy to perform in an ELISA format• Non-radioactive assay• High throughput (96 well plate format)• Assay can be completed within 5 hours

Product Details

Components:	<ul style="list-style-type: none">• 96-well Strip Microplate pre-coated with DNA probes• DNA Binding Buffer• Positive Control Sample• Specific Competitor DNA probe• Non-specific Competitor DNA probe• Assay Reagent• DTT• Wash Buffer• Primary Antibody• HRP-conjugated Secondary Antibody• TMB One-Step Substrate Reagent• Stop Solution
-------------	--

Material not included:	<ul style="list-style-type: none">• Distilled or deionized water• 100 mL and 1 liter graduated cylinders• Tubes to prepare sample dilutions Absorbent paper• Precision pipettes to deliver 2 µL to 1 mL volumes• Adjustable 1-25 mL pipettes for reagent preparation• Benchtop rocker or shaker• Microplate reader capable of measuring absorbance at 450 nm
------------------------	--

Target Details

Target:	C-JUN (JUN)
Alternative Name:	c-Jun (JUN Products)
Gene ID:	3725
UniProt:	P05412
Pathways:	MAPK Signaling , RTK Signaling , WNT Signaling , Fc-epsilon Receptor Signaling Pathway , Activation of Innate immune Response , Myometrial Relaxation and Contraction , Skeletal Muscle Fiber Development , Protein targeting to Nucleus , Toll-Like Receptors Cascades , Autophagy , Signaling of Hepatocyte Growth Factor Receptor , BCR Signaling , S100 Proteins

Application Details

Plate:	Pre-coated
Protocol:	<ol style="list-style-type: none">1. Prepare all reagents and samples as instructed in the manual.2. Add 100 µL of sample or positive control to each well.3. Incubate 2 h at RT or O/N at 4 °C.4. Add 100 µL of prepared primary antibody to each well.

Application Details

- 5. Incubate 1 h at RT.
- 6. Add 100 µL of prepared HRP-secondary antibody to each well.
- 7. Incubate 1 h at RT.
- 8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
- 9. Incubate 30 min at RT.
- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Restrictions: For Research Use only


Handling

Storage: 4 °C

Storage Comment: Upon receipt, the positive control should be removed and stored at -20° or -80°C. The remainder of the kit can be stored for up to 6 months at 2-8°C from the date of shipment. Opened Microplate Wells or reagents may be stored for up to 1 month at 2° to 8°C. Return unused wells to the pouch containing desiccant pack, reseal along entire edge. Note: The kit can be used within one year if the whole kit is stored at -20°C upon receipt. Avoid repeated freeze-thaw cycles.


Expiry Date: 6 months

Images



Activity Assay

Image 1. Transcription factor assay of c-Jun from nuclear extracts of K562 cells or K562 cells treated with PMA (50 ng/ml) for 3 hr. A. Western-blot result of c-Jun from cytoplasmic and nuclear fractions. B. Transcription factor assay of c-Jun from nuclear fractions with the TF Activity Assay Kit.



Activity Assay

Image 2. Transcription factor assay of c-Jun from nuclear extracts of K562 cells or K562 cells treated with PMA (50 ng/ml) for 3 hr with the specific competitor or non-specific competitor. The result shows specific binding of c-Jun to the conserved binding site detected by using the c-Jun TF Activity Assay Kit.