

Datasheet for ABIN5690750  
**c-MYC ELISA Kit**

2 Images



[Go to Product page](#)

## Overview

Quantity:	96 tests
Target:	c-MYC (MYC)
Reactivity:	Human
Method Type:	DNA-Binding ELISA
Application:	ELISA

## Product Details

Purpose:	Human c-Myc Transcription Factor Activity Assay. This assay uses a dsDNA coated plate with canonical c-Myc binding sequences to semi-quantitatively detect active c-Myc in lysates or nuclear extracts.
Sample Type:	Cell Lysate, Nuclear Extract
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The oligonucleotide/antibody pair provided in this kit recognizes human TFEB in whole lysates and nuclear extracts.
Characteristics:	<ul style="list-style-type: none"><li>• Specific transcription factor-DNA binding assay</li><li>• Perfect alternative to EMSA</li><li>• Easy to perform in an ELISA format</li><li>• Non-radioactive assay</li><li>• High throughput (96 well plate format)</li><li>• Assay can be completed within 5 hours</li></ul>

## Product Details

---

Components:	<ul style="list-style-type: none"><li>• 96-well Strip Microplate pre-coated with DNA probes</li><li>• DNA Binding Buffer</li><li>• Positive Control Sample</li><li>• Specific Competitor DNA probe</li><li>• Non-specific Competitor DNA probe</li><li>• Assay Reagent</li><li>• DTT</li><li>• Wash Buffer</li><li>• Primary Antibody</li><li>• HRP-conjugated Secondary Antibody</li><li>• TMB One-Step Substrate Reagent</li><li>• Stop Solution</li></ul>
-------------	--

Material not included:	<ul style="list-style-type: none"><li>• Distilled or deionized water</li><li>• 100 mL and 1 liter graduated cylinders</li><li>• Tubes to prepare sample dilutions</li><li>• Absorbent paper</li><li>• Precision pipettes to deliver 2 <math>\mu</math>L to 1 mL volumes</li><li>• Adjustable 1-25 mL pipettes for reagent preparation</li><li>• Benchtop rocker or shaker</li><li>• Microplate reader capable of measuring absorbance at 450 nm</li></ul>
------------------------	---

## Target Details

---

Target:	c-MYC (MYC)
---------	-------------

Alternative Name:	<a href="#">c-Myc (MYC Products)</a>
-------------------	--------------------------------------

Gene ID:	4609
----------	------

UniProt:	<a href="#">P01106</a>
----------	------------------------

Pathways:	<a href="#">p53 Signaling</a> , <a href="#">Cell Division Cycle</a> , <a href="#">Sensory Perception of Sound</a> , <a href="#">Transition Metal Ion Homeostasis</a> , <a href="#">Mitotic G1-G1/S Phases</a> , <a href="#">Positive Regulation of Endopeptidase Activity</a> , <a href="#">Regulation of Carbohydrate Metabolic Process</a> , <a href="#">Positive Regulation of Response to DNA Damage Stimulus</a> , <a href="#">Warburg Effect</a>
-----------	--

## Application Details

---

Plate:	Pre-coated
--------	------------

Protocol:	<ol style="list-style-type: none"><li>1. Prepare all reagents and samples as instructed in the manual.</li><li>2. Add 100 <math>\mu</math>L of sample or positive control to each well.</li><li>3. Incubate 2 h at RT or O/N at 4 <math>^{\circ}</math>C.</li></ol>
-----------	---

## Application Details

4. Add 100  $\mu$ L of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100  $\mu$ L of prepared HRP-secondary antibody to each well.
7. Incubate 1 h at RT.
8. Add 100  $\mu$ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50  $\mu$ L of Stop Solution to each well.
11. Read at 450 nm immediately.

Restrictions: For Research Use only


## Handling

Storage: 4 °C

Storage Comment: Upon receipt, the positive control should be removed and stored at -20° or -80°C. The remainder of the kit can be stored for up to 6 months at 2-8°C from the date of shipment. Opened Microplate Wells or reagents may be stored for up to 1 month at 2° to 8°C. Return unused wells to the pouch containing desiccant pack, reseal along entire edge. Note: The kit can be used within one year if the whole kit is stored at -20°C upon receipt. Avoid repeated freeze-thaw cycles.


Expiry Date: 6 months

## Images



### Activity Assay

**Image 1.** Transcription factor activity assay of TFEB from nuclear extracts of HepG2 cells or HepG2 cells treated with HBSS medium for 4 hr with the specific competitor or non-specific competitor. The result shows specific binding of TFEB to the TFEB conserved binding site.



### Activity Assay

**Image 2.** Transcription factor activity assay of TFEB from nuclear extracts of HepG2 cells or HepG2 cells treated with HBSS medium for 4 hr. After stimulation, activated TFEB is translocated into the nucleus where it binds with its corresponding DNA. A. Western-blot result of TFEB from cytoplasm and nuclear fractions. B. Transcription factor activity assay of TFEB from nuclear fractions with the TFEB Transcription Factor-Activity Assay Kit.