antibodies - online.com







PPARD ELISA Kit

Images



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	96 tests	
Target:	PPARD	
Reactivity:	Human	
Method Type:	DNA-Binding ELISA	
Application:	ELISA	
Product Details		
Purpose:	Human PPAR-delta Transcription Factor Activity Assay. This assay uses a dsDNA coated plate	
	with canonical PPAR-delta binding sequences to semi-quantitatively detect active PPAR-delta in	
	lysates or nuclear extracts.	
Sample Type:	Cell Lysate, Nuclear Extract	
Analytical Method:	Semi-Quantitative	
Detection Method:	Colorimetric	
Specificity:	The olionucleotide/antibody pair provided in this kit recognizes human PPAR-delta in whole	
	lysates and nuclear extracts.	
Characteristics:	Specific transcription factor-DNA binding assay	
	Perfect alternative to EMSA	
	Easy to perform in an ELISA format	
	Non-radioactive assay	
	High throughput (96 well plate format)	
	Assay can be completed within 5 hours	

Product Details

Components:

- 96-well Strip Microplate pre-coated with DNA probes
- · DNA Binding Buffer
- · Positive Control Sample
- Specific Competitor DNA probe
- · Non-specific Competitor DNA probe
- Assay Reagent
- DTT
- · Wash Buffer
- · Primary Antibody
- · HRP-conjugated Secondary Antibody
- TMB One-Step Substrate Reagent
- · Stop Solution

Material not included:

- Distilled or deionized water
- · 100 mL and 1 liter graduated cylinders
- · Tubes to prepare sample dilutions Absorbent paper
- Precision pipettes to deliver 2 µL to 1 mL volumes
- · Adjustable 1-25 mL pipettes for reagent preparation
- · Benchtop rocker or shaker
- · Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	PPARD	
Alternative Name:	PPAR-delta (PPARD Products)	
Gene ID:	5467	
UniProt:	Q03181	
Pathways:	Nuclear Receptor Transcription Pathway, Positive Regulation of Peptide Hormone Secretion, Steroid Hormone Mediated Signaling Pathway, Monocarboxylic Acid Catabolic Process, Smooth Muscle Cell Migration, Positive Regulation of fat Cell Differentiation	

Application Details

Plate:	Pre-coated Pre-coated
Protocol:	 Prepare all reagents and samples as instructed in the manual. Add 100 μL of sample or positive control to each well. Incubate 2 h at RT or O/N at 4 °C. Add 100 μL of prepared primary antibody to each well. Incubate 1 h at RT.

- 6. Add 100 µL of prepared HRP-secondary antibody to each well.
- 7. Incubate 1 h at RT.
- 8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
- 9. Incubate 30 min at RT.
- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Restrictions:

For Research Use only

Handling

Storage:

4°C


Storage Comment:

Upon receipt, the positive control should be removed and stored at -20° or -80°C. The remainder of the kit can be stored for up to 6 months at 2-8°C from the date of shipment. Opened Microplate Wells or reagents may be stored for up to 1 month at 2° to 8°C. Return unused wells to the pouch containing desiccant pack, reseal along entire edge. Note: The kit can be used within one year if the whole kit is stored at -20°C upon receipt. Avoid repeated freeze-thaw cycles.

Expiry Date:


6 months

Images



Activity Assay

Image 1. Transcription factor assay of PPAR-delta from 0.3 ug purified recombinate PPAR-delta protein with the specific competitor or non-specific competitor. The result shows specific binding of PPAR-delta to the PPAR-delta DNA binding site.



Activity Assay

Image 2. Transcription factor assay of PPAR-delta from purified recombinate PPAR-delta protein with PPAR-delta TF Activity Assay Kit.