

# Datasheet for ABIN5690775

# **IGF1 ELISA Kit**



### Overview

Quantity:	96 tests
Target:	IGF1
Reactivity:	Dog
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details	
Purpose:	Canine (dog) IGF-1 ELISA Kit.
Sample Type:	Cell Culture Supernatant, Cell Samples, Plasma, Serum, Tissue Lysate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA antibody pair recognizes Canine (dog) IGF-1.
Characteristics:	<ul> <li>Strip plates and additional reagents allow for use in multiple experiments</li> <li>Quantitative protein detection</li> <li>Establishes normal range</li> <li>The best products for confirmation of antibody array data</li> </ul>
Components:	<ul> <li>Pre-Coated 96-well Strip Microplate</li> <li>Wash Buffer</li> <li>Stop Solution</li> <li>Assay Diluent(s)</li> <li>Lyophilized Standard</li> <li>Biotinylated Detection Antibody</li> </ul>

### **Product Details**

- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

#### Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 μL to 1 μL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

## **Target Details**

Target:	IGF1
Alternative Name:	IGF-1 (IGF1 Products)
Gene ID:	610255
UniProt:	P33712
Pathways:	RTK Signaling, Intracellular Steroid Hormone Receptor Signaling Pathway, Peptide Hormone
	Metabolism, Hormone Activity, Regulation of Intracellular Steroid Hormone Receptor Signaling,
	Regulation of Hormone Metabolic Process, Regulation of Hormone Biosynthetic Process, Stem
	Cell Maintenance, Glycosaminoglycan Metabolic Process, Regulation of Carbohydrate
	Metabolic Process, Autophagy, Smooth Muscle Cell Migration, Activated T Cell Proliferation,
	Positive Regulation of fat Cell Differentiation

#### **Application Details**

Plate:	Pre-coated Pre-coated
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.
	2. Add 100 μL of standard or sample to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 μL of prepared biotin antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 μL of prepared Streptavidin solution to each well.
	7. Incubate 45 min at RT.
	8. Add 100 μL of TMB One-Step Substrate Reagent to each well.
	9. Incubate 30 min at RT.
	10. Add 50 μL of Stop Solution to each well.

# **Application Details**

	11. Read at 450 nm immediately.
Restrictions:	For Research Use only
Handling	
Storage:	4 °C
Expiry Date:	6 months