

# Datasheet for ABIN5690821

## **APP ELISA Kit**



### Overview

Quantity:	96 tests
Target:	APP
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details				
Purpose:	Human Amyloid-beta A4 protein/APP/PN-II ELISA Kit.			
Sample Type:	Cell Culture Supernatant, Cell Samples, Plasma, Serum, Tissue Lysate			
Analytical Method:	Quantitative			
Detection Method:	Colorimetric			
Specificity:	This ELISA antibody pair recognizes Human Amyloid-beta A4 protein.			
Characteristics:	<ul> <li>Strip plates and additional reagents allow for use in multiple experiments</li> <li>Quantitative protein detection</li> <li>Establishes normal range</li> <li>The best products for confirmation of antibody array data</li> </ul>			
Components:	<ul> <li>Pre-Coated 96-well Strip Microplate</li> <li>Wash Buffer</li> <li>Stop Solution</li> <li>Assay Diluent(s)</li> <li>Lyophilized Standard</li> <li>Biotinylated Detection Antibody</li> </ul>			

### **Product Details**

- Streptavidin-Conjugated HRP
- TMB One-Step Substrate

#### Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 μL to 1 μL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

## **Target Details**

Target:	APP		
Alternative Name:	Amyloid-beta A4 protein (APP/PN-II) (APP Products)		
Gene ID:	351		
UniProt:	P05067		
Pathways:	Caspase Cascade in Apoptosis, EGFR Signaling Pathway, Transition Metal Ion Homeostasis,		
	Skeletal Muscle Fiber Development, Toll-Like Receptors Cascades, Feeding Behaviour		

## **Application Details**

Plate:	Pre-coated
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.
	2. Add 100 μL of standard or sample to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 μL of prepared biotin antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 µL of prepared Streptavidin solution to each well.
	7. Incubate 45 min at RT.
	8. Add 100 μL of TMB One-Step Substrate Reagent to each well.
	9. Incubate 30 min at RT.
	10. Add 50 μL of Stop Solution to each well.
	11. Read at 450 nm immediately.
Restrictions:	For Research Use only

# Handling

Storage:	4 °C		
Expiry Date:	6 months		