

Datasheet for ABIN5692220

**TIM3 ELISA Kit**[Go to Product page](#)**1** Image

## Overview

Quantity:	96 tests
Target:	TIM3 (TIM 3)
Binding Specificity:	AA 22-202
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	31.2 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.2 pg/mL
Application:	ELISA

## Product Details

Purpose:	Sandwich High Sensitivity ELISA kit for Quantitative Detection of Human TIM-3/HAVCR2. 96wells/kit, with removable strips.
Brand:	PicoKine™
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Expression system for standard: NSO, Immunogen sequence: S22-G202
Sensitivity:	< 10 pg/mL
Components:	96-well plate precoated with antibody lyophilized recombinant standard

## Product Details

biotinylated antibody (dilution 1:100)  
Avidin-Biotin-Peroxidase Complex(ABC)(dilution 1:100)  
Sample diluent buffer  
Antibody diluent buffer  
ABC diluent buffer  
TMB color developing agent  
TMB stop solution  
Adhesive cover

## Target Details

Target:	TIM3 (TIM 3)
Alternative Name:	HAVCR2 ( <a href="#">TIM 3 Products</a> )
Target Type:	Virus
Background:	<p>Synonyms: Hepatitis A virus cellular receptor 2, HAVcr-2, T-cell immunoglobulin and mucin domain-containing protein 3, TIMD-3, T-cell immunoglobulin mucin receptor 3, TIM-3, T-cell membrane protein 3, HAVCR2, TIM3, TIMD3</p> <p>Tissue Specificity: Expressed in T-helper type 1 (Th1) lymphocytes. Expressed on regulatory T (Treg) cells after TCR stimulation. Expressed in dendritic cells and natural killer (NK) cells. Expressed in epithelial tissues. Expression is increased on CD4+ and CD8+ T-cells in chronic hepatitis C virus (HCV) infection. In progressive HIV-1 infection, expression is up- regulated on HIV-1-specific CD8 T-cells.</p> <p>Background: Hepatitis A virus cellular receptor 2 (HAVCR2), also known as T-cell immunoglobulin and mucin-domain containing-3 (TIM-3), is a protein that in humans is encoded by the HAVCR2 gene. The protein encoded by this gene belongs to the immunoglobulin superfamily, and TIM family of proteins. CD4-positive T helper lymphocytes can be divided into types 1 (Th1) and 2 (Th2) on the basis of their cytokine secretion patterns. Th1 cells are involved in cell-mediated immunity to intracellular pathogens and delayed-type hypersensitivity reactions, whereas, Th2 cells are involved in the control of extracellular helminthic infections and the promotion of atopic and allergic diseases. This protein is a Th1-specific cell surface protein that regulates macrophage activation, and inhibits Th1-mediated auto- and alloimmune responses, and promotes immunological tolerance.</p> <p>Cellular Localisation: Membrane, Single-pass type I membrane protein . Cell junction</p>
UniProt:	<a href="#">Q8TDQ0</a>
Pathways:	<a href="#">Regulation of Lipid Metabolism by PPARalpha</a> , <a href="#">Cancer Immune Checkpoints</a>


Application Details

Assay Time:	15 min
Plate:	Pre-coated
Restrictions:	For Research Use only

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for 6 months, at -20°C for 12 months. Avoid multiple freeze-thaw cycles(Shipped with wet ice.)
Expiry Date:	12 months

Images



ELISA

**Image 1.** Human TIM-3 PicoKine ELISA Kit standard curve