

Datasheet for ABIN5693013  
**anti-IL12B antibody (AA 23-253)**



[Go to Product page](#)

1 Image 1 Publication

## Overview

Quantity:	100 µg
Target:	IL12B
Binding Specificity:	AA 23-253
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL12B antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

## Product Details

Purpose:	Anti-IL12B Antibody Picoband®
Immunogen:	E. coli-derived human IL12B recombinant protein (Position: I23-E253).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-IL12B Antibody Picoband® (ABIN5693013). Tested in ELISA, WB applications. This antibody reacts with Human, Mouse. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.

## Target Details

Target:	IL12B
Alternative Name:	IL12B ( <a href="#">IL12B Products</a> )
Background:	<p>Synonyms: Interleukin-12 subunit beta, IL-12B, Cytotoxic lymphocyte maturation factor 40 kDa subunit, CLMF p40, IL-12 subunit p40, NK cell stimulatory factor chain 2, NKSF2, IL12B, NKSF2</p> <p>Background: Subunit beta of interleukin 12 (IL12B), also known as interleukin 12B, or interleukin-12 subunit p40, is a subunit of human interleukin 12. It is mapped to 5q33.3. This cytokine is expressed by activated macrophages that serve as an essential inducer of Th1 cells development. This cytokine has been found to be important for sustaining a sufficient number of memory/effector Th1 cells to mediate long-term protection to an intracellular pathogen. Overexpression of this gene was observed in the central nervous system of patients with multiple sclerosis (MS), suggesting a role of this cytokine in the pathogenesis of the disease.</p>
Molecular Weight:	37 kDa
Gene ID:	3593
UniProt:	<a href="#">P29460</a>
Pathways:	<a href="#">JAK-STAT Signaling</a> , <a href="#">Cellular Response to Molecule of Bacterial Origin</a> , <a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Positive Regulation of Immune Effector Process</a> , <a href="#">Activated T Cell Proliferation</a>

## Application Details

Application Notes:	<p>Western blot, 0.1-0.5 µg/mL</p> <p>ELISA, 0.1-0.5 µg/mL</p> <p>1. Entrez Gene: IL12B interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic lymphocyte maturation factor 2, p40). 2. Yokouchi, Y., Nukaga, Y., Shibasaki, M., Noguchi, E., Kimura, K., Ito, S., Nishihara, M., Yamakawa-Kobayashi, K., Takeda, K., Imoto, N., Ichikawa, K., Matsui, A., Hamaguchi, H., Arinami, T. Significant evidence for linkage of mite-sensitive childhood asthma to chromosome 5q31-q33 near the interleukin 12 B locus by a genome-wide search in Japanese families. Genomics 66: 152-160, 2000.</p>
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.


Handling

Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05 mg NaN <sub>3</sub> .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Publications

Product cited in:	Dong, Xia, Chen, Chen, Zhang: "Anti-arthritis activity of cationic materials." in: <b>Journal of cellular and molecular medicine</b> , Vol. 14, Issue 7, pp. 2015-24, (2011) ( <a href="#">PubMed</a> ).
-------------------	--

Images



Western Blotting

**Image 1.** Western blot analysis of IL12B using anti-IL12B antibody . Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each Lane was loaded with 50ug of sample under reducing conditions. Lane 1: mouse NIH3T3 cell lysates, Lane 2: human A549 cell lysates.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL12B antigen affinity purified polyclonal antibody (Catalog # ) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an

Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for IL12B at approximately 37KD. The expected band size for IL12B is at 37KD.